



FOOD AND AGRICULTURE ORGANIZATION
OF THE UNITED NATIONS

ORGANISATION DES NATIONS UNIES POUR
L'ALIMENTATION ET L'AGRICULTURE

ORGANIZACION DE LAS NACIONES UNIDAS
PARA LA AGRICULTURA Y LA ALIMENTACION

AGP: CNP/75/45
June 1975

onomy and
ysiology

Fourth Session of the FAO Technical Working Party on
Coconut Production, Protection and Processing

Kingston, Jamaica, 14-25 September 1975

IDENTIFICATION OF PREPOTENT WEST COAST TALL PALMS BASED ON PROGENY PERFORMANCE

K. Satyabalan, K.U.K. Nampoothiri and Jacob Mathew
Central Plantation Crops Research Institute
Kasaragod, Kerala, India

SUMMARY

A study of seedling characters and yield attributes of 43 open pollinated progenies of eight high yielding palms of West Coast Tall planted in 1953 has shown that the progenies of three palms are high yielding and that they are superior in spathe production and female flower production to others indicating that they are prepotents. The possibility of identifying such prepotent palms in the nursery is indicated. The progenies of such palms will help to increase the production of future plantations. Efforts to spot out more number of such palms are under way for further breeding work.

INTRODUCTION

The two important stages at which selection is practised in coconuts are at the mother palm and seedling level. In India selection of regularly yielding mother palms, yielding on an average eighty or more nuts per year (Menon and Pandalai, 1958) followed by selection of vigorous seedlings from their progeny are the usual methods followed in producing the quality planting material in coconut. But different opinions regarding the relative efficacy of mother palm selection and seedling selection have been expressed by some workers. Smith (1933) has stressed the need for seed selection as well as study of nursery results in view of the fact that certain palms persistently yield nuts of low germination and poor growth characteristics. Dwyer (1938) is of the opinion that some palms tend to produce much stronger seedlings than others. Haldane (1958) felt as a result of the findings in Ceylon reported by Pieris (1937) and others, the parent selection may be more efficient in one population and seedling selection in another. Harland (1957) has stated that all high yielding mother palms do not produce high yielding progenies and that individuals could be identified on the basis of progeny performance. He has stated that from a study of sufficiently high number of progenies from open pollinated mothers it should be possible to arrange them in order of yield and pick out mothers which in spite of having been indiscriminately pollinated by miscellaneous males are sufficiently possessed of dominant yield genes that their progeny are also high yielding.

He has compared the situation in coconut with that in cocoa where his experiments in open pollinated cocoa trees have shown that they differ in their power of transmitting high yield to their progenies. Though the genetical basis of prepotency is not clearly understood, Liyanage (1972) opined that it could be due to general combining ability. Charles (1961) evaluated that potentiality of mother palm selection and produced experimental evidence to indicate that the technique is ineffective probably because of low heritability of yield. He has recommended progeny testing as the most reliable method of detecting prepotent palms. Ninan and Pankajakshan (1961) studied the seedling characters of the open pollinated progenies and hybrids of some high yielders of West Coast Tall variety and came to the conclusion that on the basis of seedling performance it is possible to isolate high yielders which yield superior progenies from those showing inferior progeny performance.

Follow up studies of seedlings have since been made to find out how far the superior progeny performance of the open pollinated seedlings in the nursery is maintained in later years. In this paper open pollinated progenies of eight high yielding palms of the West Coast Tall variety have been studied for their seedling characters and yield performance and the results discussed with particular reference to prepotency in coconuts.

MATERIALS AND METHODS

Selected progenies raised from open pollinated nuts of eight high yielders of the West Coast Tall variety were planted in East Block (sandy loam soil) of this Institute in 1953 for a study of their performance. Data on the following characters of the forty three progenies of these high yielding palms were gathered. (1) Total number of leaves at the time of planting. (2) Girth at collar of the seedling. (3) Height of the seedling. (4) Time taken for the first flowering. (5) Number of leaves produced till 1961 since germination. (6) Number of functioning leaves in 1967. (7) Number of inflorescences produced during the five year period 1970 to 1974. (8) Number of female flowers produced during 1970 to 1974, and (9) Yield of nuts during 1970 to 1974. The number of seedlings studied in each palm varied between 4 and 10. All the seedlings were one year old at the time of planting and had received the same treatment throughout.

RESULTS AND DISCUSSION

In Table 1 are presented data on number of leaves, girth at collar and height of the seedlings at the time of planting, time taken for first flowering, mean number of leaves produced till 1961 since germination and mean number of functioning leaves on the crown in 1967. The data indicate the variations in the characters in the eight families studied. In the seedling characters studied the mean number of leaves varied from 7.0 to 8.4. The variation in the girth at collar of the seedling was from 14.3 cm to 16.7 cm and in the height of seedling from 134.3 cm to 174.4 cm. The time taken for initial flowering varied from 84.8 months (39.2.1.245) to 131.4 (29 N.312). The mean number of leaves produced till 1961 since germination indicate that families 27 N.367, 39.2.1.245, 29 S.346 and 39.2.1.205 have produced more than 70 leaves indicating their superiority in growth over others. Among the different growth characters of the seedling indicating vigour, production of leaf is the most important one. Even in adult palms it has been reported that number of leaves are significantly and positively correlated to early flowering and high yield (Patel, 1938). He has also reported that trees with a large number of leaves commence their reproductive phase earlier than trees having a lower number of leaves. In this study, progenies of families 39.2.1.245, 27 N.367, 39.2.1.205 and 29 S.346 have produced more leaves and also have more functioning leaves on their crown. They are also found to be early bearing when compared to other families.

The yield attributes of palms like number of spathes, female flower production and yield of nuts are given in Table 2. The F test indicated that for these characters, the differences between families are significant.

The range in yield of nuts and percentage of progenies yielding more than 80 nuts per palm per annum are also given. The data indicate that progenies of families 27 N.367, 29 S.346 and 39.2.1.205 are high yielding when compared to those of others. In family 27 N/367 the mean yield is 102 nuts and eighty per cent of the progenies yield more than 80 nuts. In family 29 S/346 the mean yield is 94 nuts and all the progenies yield more than 80 nuts per palm while in family 39.2.1/205 the mean yield is 85 nuts and seventy five per cent of the progenies yield more than 80 nuts per palm. In family 39.2.1/245 the leaf production as well as the spathe production are high when compared to others. The mean number of functioning leaves also is very high.

It could be seen from Table 3 that families 29 N/367, 29 S/346 and 39.2.1/205 are better than the other five families for the yield of nuts and yield attributes occupying first, second or third position, in most of the cases, when ranked. Thus based on the progeny performance these three palms can definitely be considered as prepotent palms. Such prepotent palms can be spotted out in the nursery by their progeny performance is indicated by the fact that the two families 29 S.346 and 39.2.1.205 are among the palms which were found to yield the best progenies among the 37 palms studied by Ninan and Pankajakshan (1961). The success of mother palm selection lies in the identification of such prepotent palms among the high yielders which can be identified by a study of progeny performance even in the nursery. The seedlings of such selected palms only will be able to raise the yield of coconut plantation and such vigorous selection of mother palm will go a long way to increase the production of future coconut plantations in the country. No doubt a very large number of palms will have to be screened to identify sufficient number of prepotent palms for further breeding work. It is with this objective that about 4,000 seedlings from 190 families of West Coast Tall are being studied in different locations to identify prepotent palms among them.

REFERENCES

- | | | |
|----------------------------------|------|--|
| Charles, A.E. | 1961 | Selection and breeding of the Coconut palm. <u>Trop. Agric. Trin.</u> 38: 283-96. |
| Dwyer, R.E.P. | 1938 | Coconut improvement by seed selection and plant breeding. <u>New Guinea agric. Gaz.</u> 4: 24-102. |
| Haldane, J.B.S. | 1958 | Suggestions for research in coconuts. <u>Indian Coconut J.</u> 12: 1-9. |
| Harland, S.C. | 1957 | The improvement of the coconut palm by breeding and selection. <u>Bull. 15. Coconut Res. Inst. Ceylon.</u> |
| Liyanage, D.V. | 1972 | Production of improved coconut seed by hybridisation. <u>Oleagineux.</u> 27: 597-99. |
| Menon, K.P.V. and Pandalai, K.M. | 1958 | <u>The Coconut Palm - a Monograph.</u> Indian Central Coconut Committee, Ernakulam. |

- Ninan, C.A. and
Pankajakshan, A.S. 1961 Progeny studies in coconuts.
1. Relationship between parent
yield and seedling characters of
progeny with special reference to
open pollinated and hybrid progenies
of West Coast Tall and its bearing
on the concept of prepotency in
coconuts. Indian Coconut J.
14: 100-9.
- Patel, J.S. 1938 The Coconut - A Monograph.
Govt. Press, Madras.
- Pieris, W.V.D. 1937 1. Seed selection of coconuts.
ii. Nursery management and
selection of seedlings. Trop.
Agriculturist. 88: 216-24.
- Smith, A.C. 1933 Practical seed selection of
coconuts. Malay agric. J. 21:
265-71.

Table 1

Data on the different characters of progenies of eight families

Family number and progenies studied	Seedling characters			Time taken for initial flowering till 1961 since germination	Mean number of leaves produced till 1961 since germination	Mean number of functioning leaves in 1967
	Leaves at the time of planting	Girth at the time of planting	Height at the time of planting			
	cm	cm	cm	months		
29 S.321 (10)	7.5	14.8	171.2	124.2	59.1	37.2
29 S.351 (5)	7.1	15.5	134.8	97.4	57.9	33.0
29 S.353 (4)	8.0	16.4	134.3	96.8	63.2	34.5
29 S.346 (5)	8.4 ✓	16.7 ✓	141.4	89.0 ✓	74.4	38.6 ✓
29 H.312 (5)	7.1	14.8	174.4	131.4	57.9	36.0
39.2.1.205(4)	7.5 ✓	15.8 ✓	154.0	88.0 ✓	71.3	41.0 ✓
39.2.1.245(5)	7.0 ✓	14.3 ✓	137.8	84.8 ✓	75.6	43.0 ✓
27 H.367 (5)	7.7 ✓	16.1 ✓	158.0	94.2 ✓	76.9	41.8 ✓

Ex 60.3
 7.53
 Time taken is less.
 Rate of leaf prod. mod.

1
 5
 1

29

Table 2

Yield attributes of progenies of eight families (mean of 1970 to 1974)

Family number	Number of Progenies studied	Number of spathes/tree/year	Number of female flowers produced	Yield of nuts/tree/year	Percentage of progenies yielding more than 80 nuts per palm per annum
1. 29 S.321	(10)	9.4	199.6	48.1 *(23-75)	0%
2. 29 S.351	(5)	7.6	158.6	61.8 *(45-87)	20%
3. 29 S.353	(4)	8.2	180.4	55.2 *(36-63)	0%
4. 29 S.346	(5)	11.8	269.0	84.4 *(84-109)	100% ✓✓
5. 29 N.312	(5)	10.0	208.5	64.1 *(36-95)	20%
6. 39.2.1.205	(4)	10.6	278.9	85.3 *(61-106)	75% ✓
7. 39.2.1.245	(5)	12.0	184.0	64.2 *(51-80)	20%
8. 27 N.367	(5)	10.4	319.1	101.7 *(53-143)	80% ✓✓
Gen. mean		9.9	221.6	69.1	

*Range

3 families are having more than 80 nuts & above

Table 3

Ranking of mother palms based on progeny performance

	1	2	3	4	5	6	7	8
Number of leaves	346	353	367	205	321	312	351	245
Girth at collar	346	353	367	205	351	312	321	245
Height	312	321	367	205	346	245	351	353
Time taken for initial flowering	245	205	346	367	353	351	321	312
Number of leaves produced till 1961 since germination	367	245	346	205	353	321	312	351
Number of functioning leaves in 1967	245	367	205	346	321	312	353	351
Number of spathes	245	346	205	367	312	321	353	351
Number of female flowers	367	205	346	312	321	245	353	351
Yield of nuts	367	346	205	245	312	351	353	321

Note : Block numbers of palms have been excluded for clarity of presentation.

✓