

till death. Fecundity ranges between 189–330 with an average of 249 eggs. The sex ratio in a brood works out at 1 : 3 for females to males.

The early instar caterpillars are gregarious in habit. They feed on tender foliage. The full grown caterpillars are voracious feeders and are active during night only. During the day, they congregate on tree trunks or branches and quite often escape notice. The caterpillars can completely defoliate young cashew trees.

The authors are grateful to Mr. M. C. Nambiar, Project Co-ordinator (Spices and Cashew), and Dr. K. V. Ahamed Bavappa, Director, for their keen interest in this study and to Dr. M. R. G. K. Nair, Professor of Entomology, Agricultural College and Research Institute, Vellayani, Trivandrum, Kerala State, for the valuable suggestions in the preparation of this note.

Central C. P. RADHAKRISHNAN NAIR
Plantation V. A. ABRAHAM
Crops Research G. B. PILLAI
Institute,
Kasaragod 670 124,
Kerala State,
India.

REFERENCES

- ABRAHAM, E. V. 1958. Pests of cashew (*Anacardium occidentale* L.) in South India. *Indian J. Agri. Sci.* 28 : 531–544.
- AYYAR, T. V. R. 1940. *Hand Book of Economic Entomology for South India*. Government Press, Madras, pp. 528.
- BASHEER, M. AND JAYARAJ, S. 1964. Cashewnut Pests. pp. 261–266. In *Entomology in India, Silver Jubilee Number*, *Indian J. Ent.* Entomological Society of India, New Delhi.
- FLETCHER, T. B. 1914. *Some South Indian Insects*. Government Press, Madras, pp. 565.
- SIVAGAMI, R. AND DAVID, B. V. 1968. Some insect pests of moringa (*Moringa oleifera* Linn.) in South India. *South Indian Hort.* 16 : 69–71.

Brown Rot of Turmeric

WHILE recording various pests and diseases affecting turmeric rhizomes in the germplasm collection of *Curcuma aromatica* Salisb, and *C. longa* Linn., we noticed an unusual dis-

colouration and rotting of mature rhizomes in *C. aromatica* at the time of harvest during 1973–74. Our observations are given below.

During the early stages of infection, the rhizomes are dull in colour. In advanced stages, they become deep grey to dark brown, less turgid, lose some weight, wrinkled and exhibit dry rot symptoms. The fingers are more severely affected than the mother rhizomes. When cut open affected rhizomes show dark brown necrotic lesions starting from the margin into the internal tissues. In initial stages, these necrotic lesions are localized, remain discontinuous and extend to a depth of 2–5 mm. At later stages, the lesions coalesce to form larger necrotic areas and progressively extend over a major portion of the rhizomes.

Thin hand sections of the rhizomes revealed the presence of nematodes and eggs in cortex and pith. The affected tissues were dark brown and showed, in some cases, cell disruption. When stained with cotton blue the sections showed intra and inter-cellular hyphae. From the affected tissues, *Fusarium* sp. was isolated on potato sucrose agar medium. The nematodes in the tissues were extracted, stained with lactophenol-acid fuchsin, and identified as *Pratylenchus* sp. While confirming the identification of the nematode, Dr. Orton Williams of the Commonwealth Institute of Helminthology, London, opined that it might be a new species.

The infected rhizomes were chopped and mixed with sterile soil (50 g/bag) contained in polythene bags of size 25.0 × 15.0 cm. Two pieces of seed rhizomes (about 10 g each) of *Ca 67 Jobedi* variety were sown in each bag. The plants thus raised were stunted in growth. On detailed examination after 5 months the root system was only poorly developed with varying degrees of discoloration. The rhizomes also showed brown rot symptoms. Thin hand sections showed the presence of *Pratylenchus* and *Fusarium* in the tissues. This implies a combined role for these two organisms in the causation of the malady. *Pratylenchus-Fusarium* association has been reported in the case of several diseases (Hutton, Wilkinson and Mai, 1973).

The only nematode that has been so far recorded on turmeric (*C. longa*) is the root knot nematode *Meloidogyne javanica* (Nirula and Kumar, 1963).

The authors thank Dr. Orton Williams of the Commonwealth Institute of Helminthology, London, and Dr. P. K. Koshy, Nematologist, CPCRI, Regional Station, Kayangulam for the identification of the nematode. Thanks are due to Mr. M. C. Nambiar, Project Coordinator, and Dr. K. V. Ahamed Bavappa, Director, for their encouragement.

Central Y. R. SARMA
Plantation K. K. N. NAMBIAR
Crops Research C. P. RADHAKRISHNAN NAIR
Institute,
Kasaragod 670 124,
Kerala State, India.

HUTTON, D. G., WILKINSAN, R. E., AND
MAI, W. F. 1973. Effect of two plant para-
sitic nematodes on *Fusarium* dry rot of beans.
Phytopathology 63 : 749-751.

NIRULA, K. K. AND KUMAR, R. 1963. Colla-
teral host plants of root-knot nematodes. *Curr.*
Sci. 32 : 221-222.
