

NOTE BREVI - SHORT COMMUNICATIONS

Nematology Laboratory, Central Plantation Crops Research, Institute,
Regional Station, Kayangulam, Krishnapuram 690 533, Kerala, India

RESPONSE OF SEVEN BLACK PEPPER CULTIVARS TO
MELOIDOGYNE INCOGNITA

by

P. K. KOSHY and P. SUNDARARAJU

Various authors have reported *Meloidogyne* spp. as the cause of decline and unthrifty growth of black pepper (*Piper nigrum* L.) Ayyar (1926), Nadakal (1964), D'souza *et al.* (1970) and Venkitesan (1972) in India, Ting (1975) in Malaysia, and Sharma and Loof (1974) in Brazil. In Kerala, South India, *M. incognita* (Kofoid *et* White) Chitwood is widespread and an initial inoculum of 10 larvae per plant has been found to cause severe damage to black pepper vines (Koshy *et al.*, unpublished). Seven cultivars have been screened against *M. incognita* as part of an investigation on the use of resistant pepper for nematode control.

The cultivars exposed to infection were Panniyur I, Karimunda, Kuthiravalli, Kalluvalli, Kottanadan, Narayakodi and Valiakaniakadan. Healthy cuttings of each cultivar were planted in steam sterilised sandy loam. When they had grown to 15-30 cm tall, six plants of each cultivar were transferred to pots and inoculated with approximately 5,000 freshly hatched second stage larvae of *M. incognita*. A further inoculation of 5,000 larvae to each plant was made 15 days later. The plants were maintained in a glasshouse for 7 months and then the root systems were indexed for the extent of galling by counting the number of galls in 5 cm length of root viz. mild = 1 to 5; moderate = 6 to 10; severe = 11 to 15; very severe = 16 to 20 (Table I). The roots were cut in 2 cm pieces, mixed thoroughly and a 3 g aliquot of each cultivar stained in boiling acid-fuchsin lactophenol. The stained pieces were then comminuted for 40 seconds in a Waring Blender, the volume made up to 200 ml and three 5 ml aliquots

examined for eggs, larvae and adults (Table I). Numbers of *M. incognita* in the soil were assessed from sub-samples from each pot (Table I).

All seven cultivars were susceptible in varying degrees to *M. incognita*. The highest level of infection occurred with cv. Panniyur I and the lowest with cv. Valiakaniakadan. The root galling was related to the multiplication factor. None of the cultivars could be regarded as sufficiently resistant to provide a means of control.

Table I - Response of seven black pepper cultivars to *M. incognita*.

Cultivar	Soil and root population ^a	Multiplication factor ^b	Root-gall index ^c
Panniyur - I	77,972	7.7	3.3
Karimunda	73,187	7.3	3.0
Kalluvalli	45,004	4.5	2.0
Kuthiravalli	43,705	4.3	1.8
Naranyakodi	22,965	2.2	1.7
Kottanadan	16,863	1.6	1.3
Valiakaniakadan	9,600	1.0	1.1
S.E.	13,165		
M.S.D.	36,205		

a) Mean of 6 replicates (Total/pot); b) Soil and root population increase; c) See text.

The authors thank Dr. K. V. Ahamed Bavappa, former Director, C.P.C.R.I., Kasaragod for providing necessary facilities and Shri M. C. Nambiar, Project Co-ordinator, Cashew and Spices for encouragement.

L I T E R A T U R E C I T E D

- AYYAR P. N. K., 1926 - A preliminary note on the root gall nematode, *Heterodera radicicola* and its economic importance in South India. *Madras Agric. J.*, 14: 113-118.
- D'SOUZA G. I., KUMAR A. C., KASIVISWANATHAN P. R. and SHAMANNA H. V., 1970 - Relative distribution and prevalence of plant parasitic nematodes in the coffee tracts of South Western India. *Indian Coffee*, 34: 329-342.
- KOSHY P. K., SOSAMMA V. K. and SUNDARARAJU P., 1977 - Screening of plants used as pepper standards against root-knot nematode. *Indian Phytopath.*, 30: 128-129.

- NADAKAL A.M. 1964 - Studies on plant parasitic nematodes of Kerala III. An additional list of plants attacked by root-knot nematode *Meloidogyne* sp. *J. Bombay nat. His. Soc.*, 61: 467-469.
- SHARMA R.D. and LOOF P.A.A., 1974 - Nematodes of the cocoa region of Bahia, Brazil IV. Nematodes in the rhizosphere of pepper (*Piper nigrum* L.) and clove (*Eugenia caryophyllata* Thumb.). *Rev. Theobroma*, 4: 26-32.
- TING W.P., 1975 - Plant pathology in peninsular Malaysia. *Rev. Pl. Path.*, 54: 287-305.
- VENKITESAN T.S., 1972 - On the occurrence of plant parasitic nematodes associated with different crops in Cannanore District, Kerala. *Agric. Res. J. Kerala*, 10: 179-180.

Accepted for publication on 16 November, 1978.