



Two new alkaloids isolated from traditional Chinese medicine Binglang the fruit of *Areca catechu*

Mengru Cao^{a,1}, Hanwen Yuan^{a,1}, Muhammad Daniyal^a, Huanghe Yu^a, Qingling Xie^a, Yingkai Liu^a, Bin Li^a, Yuqing Jian^a, Caiyun Peng^a, Dianbo Tan^{b,*}, Yanmei Peng^b, M. Iqbal Choudhary^a, Atta-ur Rahman^a, Wei Wang^{a,*}

^a TCM and Ethnomedicine Innovation & Development International Laboratory, Academician Atta-ur-Rahman Belt and Road Traditional Medicine Research Center, School of Pharmacy, Hunan University of Chinese Medicine, Changsha, Hunan 410208, PR China

^b Institute of Chinese Medicine, Hunan Academy of Chinese medicine, Changsha, Hunan 410208, PR China

ARTICLE INFO

Keywords:

Areca catechu

Areaceae

Alkaloids

Binglang

Structural identification

TCM

ABSTRACT

Binglang, the fruit of *Areca catechu* L, has a long history as an important Chinese herbal medicine. Two new alkaloids (**1** and **2**), along with forty-one known compounds (**3–43**) were isolated from the dried fruit of *Areca catechu* L. The structures were elucidated on basis of the IR, UV, MS and 1D, 2D NMR spectroscopic data. Compounds **26** and **33** showed weak cytotoxicity against human gastric cancer cell line (BGC-823) with IC₅₀ of 15.91 μM and 20.13 μM, respectively.

1. Introduction

Areca catechu L, called “Binglang” in Chinese, belongs to the family Areaceae which is comprised of fifteen species. This plant is widely distributed in southern and southeast Asia including China, India, Pakistan, Indonesia, Malaysia, Philippines and New Guinea. In Hainan province of China for its important economic and medicinal benefits [1,2]. The fruits of *A. catechu* (frequently known as areca nut) are popular chewable items in Asian countries [3,4]. In addition, the peel and seed of *A. catechu* have been used as traditional Chinese medicine (TCM) since 1700 years [5]. Since 1953, the areca seed have been listed in the Pharmacopoeia of the People's Republic of China and over 100 prescriptions containing areca seed such as Binglang sixiao pill, Simo decoction, and Jianweixiaoshi tablets have been utilized for treatment of parasitic diseases, abdominal distension, abdominal pain, and jaundice [6]. Dafupi, which is the peel of *A. catechu*, is used as TCM to treat ascites and edema [6,7].

The major chemical constituents in *A. catechu* have earlier been identified are; alkaloids, flavonoids, tannins, triterpenes, and fatty acids [2,8]. A number of pharmacological activities such as anti-parasitic effect, anti-depressive effect, anti-fatigue effect, anti-oxidant effect and increasing gastrointestinal motility properties [9,10]. The alkaloids of *A. catechu* were found to have anti-taeniasis, anti-tumor, antibacterial,

and antifungal effects [8]. Chemical constituents of *A. catechu*, including catechin, chrysoeriol, isorhamnetin, luteolin, isovanillic acid, protocathechuic acid, showed significant antioxidant activity [2,4,8]. These pharmacological activities indicated that the *A. catechu* have a promising future for treating diseases, particular parasitosis, digestive system diseases and nervous system disorders.

In order to find structurally unique and bioactive natural products, 95% EtOH extract of this plant were separated, which led to the isolation of two new compounds, named as acatechu A (**1**) and acatechu B (**2**) (Fig. 1), along with forty-one known ones (**3–43**). Compounds **7–11**, **13–14**, **16–17**, **23–27**, **34–35**, **40**, and **42–43** were isolated from this plant for the first time. In addition, all isolated compounds were evaluated for their cytotoxic activity against four human cancer cell lines (Hela, BGC-823, Hep G2 and HCT-116). Compounds **26** and **33** showed weak cytotoxicity against human gastric cancer cell line BGC-823.

2. Results and discussion

Compound **1** was obtained as yellowish power. The molecular formula of **1** was C₂₀H₂₇NO₇ determined by its negative HR-ESI-MS *m/z* 416.1673 [M-Na][−] (calcd for C₂₀H₂₇NO₇[−], 416.1685). The UV spectrum of **1** showed the absorption maxima at 206 and 279 nm. The IR spectrum indicated the presence of hydroxyl (3420 cm^{−1}), carbonyl

* Corresponding authors.

E-mail addresses: 908191609@qq.com (D. Tan), wangwei402@hotmail.com (W. Wang).

¹ These authors contributed equally to this work.




Fig. 1. The chemical structures of 1–2.




Fig. 2. The key ^1H - ^1H COSY (—) and HMBC (—) correlations of 1–2.

(1717 cm^{-1}), and aromatic ring (1576 and 1463 cm^{-1}). The ^1H NMR spectrum showed four aromatic proton signals [δ_{H} 7.54 (1H, d, $J = 8.3$, H-7), 7.52 (1H, d, $J = 7.9$, H-4), 7.19 (1H, ddd, $J = 8.2$, 7.2, 1.1, H-6), 7.09 (1H, m, H-5)]. The proton signals in the region of 5.44 ppm–3.50 ppm [δ_{H} 5.44 (1H, d, $J = 9.1$, H-1'), 3.92 (1H, m, H-2'), 3.60 (1H, t, $J = 9.0$, H-3'), 3.57 (1H, dd, $J = 5.7$, 2.2, H-5'), 3.50 (1H, m, H-4'), 3.87 (1H, d, $J = 2.0$, H-6'), 3.71 (1H, dd, $J = 12.2$, 5.7, H-2-6')] suggested the presence of a sugar moiety, which was identified as glucopyranoside based on analysis of HMBC and ^1H - ^1H COSY NMR data (Fig. 2). In ^{13}C NMR spectrum signals for a carbonyl groups [δ_{C} 174.3C-9], a benzene ring [δ_{C} 138.4C-7a, 129.7C-3a, 123.1C-6, 120.9C-5, 119.8C-4, 111.6C-7], and a glucopyranoside unit [δ_{C} 86.7C-1', 80.6C-3', 78.9C-5', 73.8C-2', 71.5C-4', 62.8C-6'] were observed. The ^1H and ^{13}C NMR spectra of 1 were closely similar to those of 3-carboxymethyl indole 1- N - β -D-glucopyranoside [11], with the only difference of an extra signal for *n*-butyl group signals [δ_{H} 4.12 (2H, t, $J = 6.6$, H-1''), 1.62 (2H, m, H-2''), 1.39 (2H, dq, $J = 14.8$, 7.4, H-3''), 0.93 (3H, t, $J = 7.4$, H-4''), δ_{C} 65.9C-1'', 31.9 C-2'', 20.2C-3'', 14.1 C-4''] for compound 1, which was placed at C-9 based on HMBC correlation from H-1'' [δ_{H} 4.12 (2H, t, $J = 6.6$)] to C-9 [δ_{C} 174.3] (Fig. 2). The β -D-configuration of the glucopyranoside was determined by comparison with standard D-glucopyranoside using HPLC method after acid hydrolysis, and the coupling constant shift of the anomeric proton [δ_{H} 5.44 (d, $J = 9.1$)] [12]. Thus, compound 1 was elucidated as 3-carboxymethylbutyl ester indole 1- N - β -D-glucopyranoside, and named as acatechu A accordingly.

Compound 2 was obtained as yellowish powder. The molecular formula of 2 was determined as $\text{C}_{22}\text{H}_{27}\text{NO}_9$ by negative HR-ESI-MS at m/z 466.1737 [$\text{M} + \text{H}_2\text{O}-\text{H}^-$] (calcd for $\text{C}_{22}\text{H}_{27}\text{NO}_9^-$, 466.1713). The UV spectrum of 2 showed the absorption maxima at 204 and 285 nm. The IR spectrum implicated the presence of hydroxyl (3375 cm^{-1}), carbonyl (1733 cm^{-1}), and aromatic ring (1513 and 1447 cm^{-1}). The ^1H NMR spectrum signals in the region of 7.34–6.58 ppm [δ_{H} 7.34 (2H, m, H-2'' 6''), 7.31 (2H, d, $J = 6.8$, H-3'' 5''), 7.27 (1H, t, $J = 7.1$, H-4''), 6.80 (1H, d, $J = 2.7$, H-2), 6.69 (1H, d, $J = 8.6$, H-5), 6.58 (1H, dd, $J =$

8.6, 2.7, H-6)] indicated the presence of two benzene rings. In addition, the signals of a glucopyranoside unit [δ_{H} 4.74 (1H, d, $J = 7.5$, H-1'), 3.90 (1H, dd, $J = 12.0$, 2.2, H-6'), 3.68 (1H, dd, $J = 12.0$, 5.9, H-2-6'), 3.41 (1H, m, H-3'), 3.42 (1H, m, H-4'), 3.39 (1H, dd, $J = 5.9$, 2.3, H-2'), 3.36 (1H, d, $J = 8.4$, H-5')] were observed in the ^1H NMR spectrum of 2. Two benzene rings signals [δ_{C} 152.8 C-1, 149.3 C-3, 142.9 C-4, 137.3 C-1'', 130.4C-3'' 5'', 129.9C-2'' 6'', 128.4C-4'', 115.9C-5, 109.9C-6, 103.8C-2], and a glucopyranoside unit signals [δ_{C} 103.8C-1', 74.9C-2', 78.0C-3', 78.2C-4', 71.6C-5', 62.7C-6'] were also observed in the ^{13}C NMR spectrum. Comparison of the ^1H and ^{13}C NMR data of 2 with those of 2,4-dihydroxyphenyl 6- O -(3-hydroxy-1-oxo-3-phenyl-propyl)- β -D-glucopyranoside [13] suggested that these two compounds were structurally similar. However, two hydrogens in 2 were attached to C-2 and C-9'', instead of two hydroxyl groups in the known compound, respectively. In addition, a methoxy group [3.83 (3H, s)] was attached to C-3 based on HMBC correlations of $\text{H}_3\text{-OCH}_3$ [δ_{H} 3.83 (3H, s)] to C-3 [δ_{C} 149.3] (Fig. 2), and an amino-group attached to C-8'' based on downfield shift of C-8'' [δ_{H} 3.78 (1H, dd, $J = 8.9$, 4.3), δ_{C} 57.58] [14]. Acid hydrolysis of compound 2 yielded a D-glucose which was identified by comparison with standard D-glucopyranoside using HPLC method. The anomeric proton [δ_{H} 4.74 (d, $J = 7.5$)] indicated the β -configuration for the glucopyranosyl moiety [12]. Thus, compound 2 was identified to be 3-methoxy-4-hydroxyphenyl 6- O -(2-amino-1-oxo-3-phenyl-propyl)- β -D-glucopyranoside, which was named as acatechu B.

In addition, forty-one known compounds including arecoline (3) [15], areca (4) [15], methyl nicotinate (5) [16], ethyl nicotinate (6) [16], β -D-fructopyranoside, ethyl (7) [17], β -D-fructopyranoside, butyl (8) [17], butyl quinate (9) [18], 2,5-dihydro-2-hydroxy-5-oxo-2-furancarboxylic acid (10) [19], β -D-fructofuranoside, butyl (11) [16], resveratrol (12) [20], 2-(4-hydroxy-phenyl) ethyl- β -D-glucopyranoside (13) [21], stroside B (14) [22], fumaric acid (15) [23], β -D-glucopyranoside, benzyl (16) [24], 5-methyl-2-(1-methylethyl)-cyclohexanol (17) [25], 2-hydroxy-5-methoxy-benzoic acid (18) [26], 4-hydroxy-benzoic acid (19) [27], 4-hydroxy-3-methoxy-benzoic acid (20) [17], 3-hydroxy-4-methoxy-benzoic acid (21) [27], 3,4-dihydroxy-benzoic acid (22) [28], 2-hydroxyethyl-5-methoxyphenyl- β -D-glucopyranoside (23) [29], 2-(β -D-glucopyranosyloxy)-5-methoxy-benzoic acid (24) [30], 3,4-dimethoxyphenyl- β -D-glucopyranoside (25) [31], 2,4-dimethoxyphenyl- β -D-glucopyranoside (26) [32], 4-hydroxy-3-methoxyphenyl- β -D-glucopyranoside (27) [33], (+)-catechin (28) [34], (–)-catechin (29) [35], liquiritigenin (30) [36], quercetin (31) [37], cyanidenon (32) [28], isorhamnetin (33) [38], isorhamnetin 3- O -(6''- O - α -L-rhamnopyranosyl)- β -D-glucopyranoside (34) [39], methyl rhamnoside (35) [40], 6- O -methylemodin (36) [41], chrysophanol (37) [41], proanthocyanidin B₁ (38) [42], β -sitosterol (39) [43], stigmastane-3 β ,5,6 α -triol (40) [25], arborinol (41) [44], β -carotenoid (42) [45], and ginseng triol saponin Rg 1 (43) [46] were isolated and identified. Their structures were determined by comparing the spectroscopic data with those

reported in literature. The chemical structures and NMR spectra of all known compounds is presented in the Supplementary material.

All isolated compounds were evaluated for their cytotoxic activities against four human cancer cell lines (Hela, BGC-823, Hep G2 and HCT-116). Compounds **26** and **34** showed weak cytotoxicity against BGC-823 cell line with IC₅₀ of 15.91 μ M and 20.13 μ M, respectively.

3. Experimental

3.1. General experimental procedures

The HR-ESI-MS spectra were performed on Waters UHPLC-H-CLASS/XEVO G2-XS Qtof. NMR data were recorded on Bruker AV-600 spectrometers with TMS as an internal standard. Column chromatographic separations were carried out with silica gel, (200–300 mesh and 300–400 mesh), octadecylsilyl (ODS) (Qingdao Marine Chemical Inc., P. R. China) and Sephadex LH-20 (Sigma-Aldrich). Semipreparative HPLC was performed on an Agilent 1206 high performance liquid chromatograph with a Agilent C₁₈ (34 mm $\times$ 25 cm) column. Fractions were monitored by TLC (Qingdao Marine Chemical Inc., P. R. China) and spots were visualized by heating silica gel plates sprayed with vanillin in 5% H₂SO₄ solution. Microplate reader (EnSpire 2300, PerkinElmer) and CO₂ incubator (New Brunswick) were used in cytotoxicity bioassays.

3.2. Plant material

The dried fruits of *Areca catechu* L. were collected in 2016 from Hainan Province of China. The plant was identified by Prof. Li-min Gong (School of Pharmacy, Hunan University of Chinese Medicine). A voucher specimen (No. 201610) was deposited at TCM and Ethnomedicine Innovation & Development International Laboratory, Innovative Materia Medica Research Institute, Hunan University of Chinese Medicine, Changsha, P. R. China.

3.3. Extraction and isolation

The dried fruits of *A. catechu* (50 Kg) were extracted twice with 1000 L of 95% ethanol for 2 h under reflux extraction. And all the solvents were evaporated under vacuum to obtain crude extract (1.2 Kg). The obtained extract was suspended in water (1500 mL) and partitioned with PE (2 L $\times$ 3 times), CH₂Cl₂ (2 L $\times$ 3 times), EtOAc (2 L $\times$ 3 times) and *n*-BuOH (2 L $\times$ 3 times) to get four fractions after removing the solvent under vacuum, respectively.

The PE (197.2 g) and CH₂Cl₂ (7.2 g) fractions were combined and then subjected to silica gel column chromatography (CC) (2000 g, 16 $\times$ 35 cm) eluted stepwise with PE-EtOAc system. The collected fractions were combined based on TLC analysis to afford 20 main fractions. These fractions were further separated by silica gel CC using gradient system (PE-EtOAc & CH₂Cl₂-MeOH) to afford **3–6**, **17–21**, and **39–41**.

The EtOAc fraction (34.9 g) was separated by silica gel CC to yield fractions A-E. The subfraction B was purified by silica gel CC (3 $\times$ 45 cm; eluted with PE-EtOAc) to afford **12** and **15**. The subfraction D was subjected to column chromatography over sephadex LX-20 (eluting with CHCl₃:MeOH = 1:1) to yield **28–33**, and **36–38**.

The *n*-butanol fraction (57 g) was separated stepwise on silica gel CC (1500 g, 10 $\times$ 32 cm) eluted with CHCl₃-MeOH-H₂O system (10:1:0.1, 8:2:0.1, 7:3:0.1, 6:4:0.1, 3:7:0.1, 0:1:0, v:v:v) by increasing polarity, to afford 12 main fractions (Fr. 1 ~ Fr. 12). Fr. 2 (2.6 g) was subjected to silica gel CC (2.5 $\times$ 62 cm, 70 g) using gradient system (CHCl₃-MeOH-H₂O) to afford **7–11** and thirteen fractions (A₁ ~ A₁₃). Compound **1** (4 mg) was obtained from A₆ by ODS column chromatography (2.5 $\times$ 8.5 cm; eluted with MeOH-H₂O). Fr. 5 (600 mg) was purified by sephadex LX-20 CC (2.5 $\times$ 96 cm) eluting with CHCl₃-MeOH system to yield compound **2** (3 mg) and eleven subfractions (B₁ ~ B₁₁). The

Table 1
¹H and ¹³C NMR data (δ in ppm) of **1** (150 and 600 MHz, MeOD).

Position	δ_c	δ_H mult (J in Hz)
2	125.4	7.37 (1H, s)
3	109.8	–
3a	129.7	–
4	119.8	7.52 (1H, d, 7.9)
5	120.9	7.09 (1H, m)
6	123.1	7.19 (1H, ddd, 8.2, 7.2, 1.1)
7	111.6	7.54 (1H, d, 8.3)
7a	138.4	–
8	31.9	3.78 (2H, s)
9	174.3	–
1'	86.7	5.44 (1H, d, 9.1)
2'	73.8	3.92 (1H, m)
3'	80.6	3.60 (1H, t, 9.0)
4'	71.5	3.50 (1H, m)
5'	78.9	3.57 (1H, dd, 5.7, 2.2)
6'	62.8	3.87 (1H, d, 2.0)
		3.71 (1H, dd, 12.2, 5.7)
1''	65.9	4.12 (2H, t, 6.6)
2''	31.8	1.62 (2H, m)
3''	20.2	1.39 (2H, dq, 14.8, 7.4)
4''	14.0	0.93 (3H, t, 7.4)

subfraction B₆ was purified by ODS column chromatography (2.5 $\times$ 8.5 cm; eluted with MeOH-H₂O) to afford **22** and **34–35**. Fr 3 and Fr 4 (2.5 g) were separated by chromatography column on silica gel (2.5 $\times$ 52 cm, 80 g) to afford **23–27** and **44–45**. Fr. 6 (2.1 g) was purified by ODS column chromatography (3.5 $\times$ 15 cm; eluted with MeOH-H₂O) to yield **16**, **13–14**, and **42–43**.

3.3.1. *Acatechu A* (1)

Yellowish powder; [α]_D24.5 D + 0.3 (c 0.2, MeOH); UV (MeOH) λ_{max} (log ϵ) 206 nm (4.19) and 279 nm (3.42); IR (ART) ν_{max} 3420, 1717, 1576, 1463 cm^{−1}; negative ion HR-ESI-MS m/z 416.1673 [M-Na][−] (calcd for C₂₀H₂₇NO₇[−], 416.1685); ¹H NMR (600 MHz, MeOD) and ¹³C NMR (150 MHz, MeOD) data were showed in Table 1.

3.3.2. *Acatechu B* (2)

Yellowish powder; [α]_D24 D + 0.15 (c 0.4, MeOH); UV (MeOH) λ_{max} (log ϵ) 204 nm (4.81) and 285 nm (3.81); IR (ART) ν_{max} 3375, 1733, 1615, 1513, 1447 cm^{−1}; negative ion HR-ESI-MS m/z 466.1737 [M + H₂O-H][−] (calcd for C₂₂H₂₇NO₉[−], 466.1713); ¹H NMR (600 MHz, MeOD) and ¹³C NMR (150 MHz, MeOD) data were showed in Table 2.

3.4. Cytotoxicity bioassays

MTT assays were performed to evaluate the cytotoxicity activities against Hela, BGC-823, Hep G 2, and HCT-116 cell lines. Briefly, all cells were seeded in 96-well plates at a density of about 5 $\times$ 10⁴ cells per well, respectively. Each cancer cell line was exposed to the test compound, which was dissolved in DMSO, at various concentrations for 48 h, with paclitaxel as positive control. After replaced the cell culture with cell medium, 100 μ L MTT solution (0.5 mg/mL in DMEM culture medium) was added to each well for additional 4 h incubation. The cell medium containing MTT was then replaced with 150 μ L DMSO. Afterwards, the micro-plate was incubated at 37 °C for 5 min and then shaken at room temperature for 15 min for complete dissolution of formazan. Finally, the absorbance of each well at 592 nm was recorded by microplate reader (EnSpire 2300) to determine the relative cell viability [47,48].

In conclusion, two new alkaloids (**1** and **2**), and forty-one known compounds (**3–43**) were isolated from the dried fruit of *A. catechu*. The present study is the first report of phytochemical investigation on the *n*-BuOH of *A. catechu* L. Different from previous reported alkaloids of *A. catechu*, two new alkaloids both were linked with glucopyranoside

Table 2
¹H and ¹³C NMR data (δ in ppm) of **2** (150 and 600 MHz, MeOD).

Position	δ _C	δ _H mult (J in Hz)
1	152.8	–
2	103.8	6.80 (1H, d, 2.7)
3	149.3	–
4	142.9	–
5	115.9	6.69 (1H, d, 8.6)
6	109.9	6.58 (1H, dd, 8.6, 2.7)
1'	103.8	4.74 (1H, d, 7.5),
2'	74.9	3.39 (1H, dd, 5.9, 2.3)
3'	78.0	3.41 (1H, m)
4'	78.2	3.42 (1H, m)
5'	71.6	3.36 (1H, d, 8.4)
6'	62.7	3.90 (1H, dd, 12.0, 2.2) 3.68 (1H, dd, 12.0, 5.9)
1''	137.3	–
2'', 6''	129.9	7.34 (2H, m)
3'', 5''	130.4	7.31 (2H, d, 6.8)
4''	128.4	7.27 (1H, t, 7.1)
7''	174.1	–
8''	57.6	3.78 (1H, dd, 8.9, 4.3)
9	38.3	3.34 (1H, d, 3.8)
		3.00 (1H, dd, 14.5, 8.9)
3-OCH ₃	56.4	3.83 (3H, s)

moiety. Compound **1**, an indole alkaloid, was isolated from Arecaceae family for the first time. The indole *N*-glycoside derivatives are the regulator of plant growth and development, and analogs of **1** have been isolated or synthesized previously [11,49]. For compound **2**, it is rare that the phenylalanine group attached to C-6' of glucopyranoside moiety in natural products. In addition, all isolated compounds were evaluated for their cytotoxic activities against four human cancer cell lines (Hela, BGC-823, Hep G2 and HCT-116). Compounds **26** and **33** showed weak cytotoxicity against human gastric cancer cell line BGC-823 with IC₅₀ of 15.91 μM and 20.13 μM, respectively.

Funding

This work was supported by the National Key R&D Program of China (2018YFC1707900), National Natural Science Foundation of China (81874369, 81803708, 81673579), Hunan Provincial Natural Science Foundation (2018JJ2293), Key Research and Development Programs of Hunan Science and Technology Department (2018SK2113, 2018SK2119, 2018WK2081).

Acknowledgments

The authors are appreciative to Hunan Province Betel-Nut Association and Dr. Bin Liu from Hunan Province Key Laboratory of Plant Functional Genomics and Developmental Regulation, College of Biology, Hunan University.

Declaration of Competing Interest

No potential conflict of interest was reported by the authors.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fitote.2019.104276>.

References

- [1] C.D. Heatubun, J. Dransfield, T. Flynn, S.S. Tjitroedirdjo, J.P. Moge, W.J. Baker, A monograph of the betel nut palms (Arecaceae) of East Malesia, Bot. J. Linn. Soc. 168 (2012) 147–173.
- [2] M.S. Amudhan, V.H. Begum, K.B. Hebbar, A review on phytochemical and pharmacological potential of *Arecaceae* L. seed, Int. J. Pharm. Sci. Res. 3 (2012) 4151–4157.
- [3] A.A. Franke, L. Biggs, J.Y. Yew, J.F. Lai, Areca alkaloids measured from buccal cells using DART-MS serve as accurate biomarkers for areca nut chewing, Drug Test Anal. 3 (2019) 1942–1960.
- [4] D.Y. Lim, I.S. Kim, Arecoline inhibits catecholamine release from perfused rat adrenal gland, Acta Pharm. Sin. 27 (2006) 71–79.
- [5] Chinese Material Medica, Science and Technology Press, Shanghai, 1999.
- [6] Chinese Pharmacopoeia Commission, Chinese Pharmacopoeia, Chemical Industry Press, Beijing, 2015 (Chapter 1).
- [7] Dictionary of Chinese Materia Medica, Science and Technology Press, Shanghai, 1977.
- [8] W. Peng, Y.J. Liu, N. Wu, T. Sun, X.Y. He, Y.X. Gao, C.J. Wu, Areca catechu L. (Arecaceae): a review of its traditional uses, botany, phytochemistry, pharmacology and toxicology, J. Ethnopharmacol. 164 (2015) 340–356.
- [9] A.H. Gilani, M.N. Ghayur, Z.S. Saify, S.P. Ahmed, M.I. Choudhary, A. Khalid, Presence of cholinomimetic and acetylcholinesterase inhibitory constituents in betel nut, Life Sci. 75 (2004) 2377–2389.
- [10] P. Boniface, S.K. Verma, H.S. Cheema, Evaluation of antimalarial and antimicrobial activities of extract and fractions from *Arecaceae*, Inter J. Infect. Dis. 21 (2014) 228–229.
- [11] B. Schwarz, T. Hofmann, Isolation, structure determination, and sensory activity of mouth-drying and astringent nitrogen-containing phytochemicals isolated from red currants (*Ribes rubrum*), J. Agric. Food Chem. 55 (2007) 1405–1410.
- [12] M. Li, L. Bin, X. Ziming, T. Ying, Z. Dan, W.J. Rui, J.X. Dong, F.J. Xiao, Anticancer effects of five biflavonoids from Ginkgo Boloba. Male flowers in vitro, Molecules. 24 (2019) 1496–1509.
- [13] L. Verotta, F. Orsini, G. Pelizzoni, C.B.R. Torri, Polyphenolic glycosides from african proteaceae, J. Nat. Prod. 62 (1999) 1526–1531.
- [14] J.P. Tian, Y.W. Yin, H.B. Sun, X.F. Luo, Magnesium chloride: an efficient ¹³C NMR relaxation agent for amino acids and some carboxylic acids, J. Magnetic Reson. 159 (2002) 137–144.
- [15] P. Srinivasan, R.L. Lichter, ¹³C NMR spectral studies of arecoline, hordenine, strychnine and brucine, Orga Magnetic Reso. 8 (1976) 198–201.
- [16] D.P. Bora, B. Deb, A.L. Fuller, A.Z.S. Alexandra, J.D. Woollins, Dicarboxyliridium(II) complexes of pyridine ester ligands and their reactivity towards various electrophiles, Inorg. Chim. Acta 363 (2010) 1539–1546.
- [17] S.Y. Lee, S.U. Choi, J.H. Lee, D.U. Lee, K.R. Lee, A new phenylpropane glycoside from the rhizome of Sparganium stoloniferum, Arch. Pharm. Res. 33 (2010) 515–521.
- [18] Y.B. Yang, X. Li, Q. Yang, Z.J. Wu, L.N. Sun, Study on chemical constituents of papaya rugosa, Acad. J. Sec. Med. Univ. (2009) 1195–1198.
- [19] X.J. Li, J.W. Cubbage, T.A. Tetzlaff, S.J. William, Photocatalytic degradation of 4-Chlorophenol. 1. The hydroquinone pathway, J. Organomet. Chem. 64 (1999) 8509–8524.
- [20] D.H. Lee, K.P.L. Bhat, H.H.S. Fong, N.R. Farnsworth, J.M. Pezzuto, A.D. Kinghorn, Aromatase inhibitors from Broussonetia papyrifera, J. Nat. Prod. 64 (2001) 1286–1293.
- [21] J. Landtag, A. Baumert, T. Degenkolb, J. Schmidt, V. Wray, D. Scheel, D. Strack, D. Rosahl, Accumulation of tyrosol glucoside in transgenic potato plants expressing a parsley tyrosine decarboxylase, Phytochemistry. 60 (2002) 683–689.
- [22] K.W. Woo, W.S. Suh, L. Subedi, S.Y. Kim, S.U. Choi, K.H. Kim, K.R. Lee, Phenolic derivatives from the stems of Lagerstroemia indica and their biological activity, Heterocycles. 91 (2015) 2355–2367.
- [23] S. Prachayasittikul, S. Suphamong, A. Worachartcheewan, R. Laung, S. Ruchirawat, V. Prachayasittikul, Bioactive metabolites from Spilanthes acmella Murr, Molecules. 14 (2009) 850–867.
- [24] D.S. Seigler, G.F. Paulib, A. Nahrstedt, R. Leen, Cyanogenic allosides and glucosides from Passiflora edulis and Carica papaya, Phytochemistry. 60 (2002) 873–882.
- [25] T. Mino, T. Hasegawa, Y. Shirai, M. Sakamoto, T. Fujita, N,O-ligand accelerated zinc-catalyzed transesterification of alcohols with vinyl esters, J. Organomet. Chem. 692 (2007) 4389–4396.
- [26] Y.H. Zhang, J.Q. Yu, Pd(II)-catalyzed hydroxylation of Arenes with 1 atm of O₂ or air, J. Amer. Chem. Soc. 131 (2009) 14654–14655.
- [27] H.Y. Ding, H.C. Lin, C.M. Teng, Y.C. Wu, Phytochemical and pharmacological studies on Chinese Paeonia species, J. Chin. Chem. Soc. 47 (2000) 381–388.
- [28] G. Flamini, E. Antognoli, I. Morelli, Two flavonoids and other compounds from the aerial parts of Centaurea bracteata from Italy, Phytochemistry. 57 (2001) 559–564.
- [29] M.G. Ghazooly, A.M. Lakany, A.S. Mohamed, H.A. Amal, Chemical constituents of Helichrysum conglobatum growing in Egypt, Nat. Prod. Sci. 9 (2003) 213–219.
- [30] D.J. Zuchowski, L.K. Pecio, R. Emilia, S. Anna, New phenolic compounds from the roots of lentil (Lens culinaris), Helv. Chim. Acta. 99 (2016) 674–680.
- [31] C. Arevalo, I. Ruiza, A.L. Piccinelli, L. Campone, L. Rastrelli, Phenolic derivatives from the leaves of Martinella obovata (Bignoniaceae), Nat. Prod. Commun. 6 (2011) 957–960.
- [32] H.J. Kwak, S. Park, N.Y. Kim, G.J. Yoo, J.H. Park, Y. Oh, N.X. Nhim, S.H. Kim, Neuraminidase inhibitory activity by compounds isolated from aerial parts of Rhinacanthus nasutus, Nat. Prod. Res. 32 (2018) 2111–2115.
- [33] X.M. Zhou, C.J. Zheng, X.P. Song, G.Y. Chuen, Y.H. Cao, A.Q. Huang, S.H. Su, W.X. Chen, H. Yang, A new phenolic glycoside from Saprosmia merrillii, Nat. Prod. Res. 21 (2016) 2429–2433.
- [34] K. Yoshiki, N. Genichiro, N. Itsuo, Tannins and related compounds. XLV. Rhubarb. (5). isolation and characterization of flavan-3-ol and procyanidin glucosides, Chem. Pharm. Bull. 4 (1986) 08–22.
- [35] M.B. Zhang, G.E. Jagdmann, Z.M. Van, R. Sheeler, P. Beckett, H. Schroeter, Chemical synthesis and characterization of Epicatechin Glucuronides and sulfates:

- bioanalytical standards for Epicatechin metabolite identification, *J. Nat. Prod.* 76 (2013) 157–169.
- [36] R. Wen, H. Lv, Y. Jiang, P.F. Tu, Anti-inflammatory Flavanones and Flavanols from the roots of *Pongamia pinnata*, *Planta Med.* 84 (2018) 1174–1182.
- [37] T. Fossen, A.T. Pedersen, O.M. Andersen, Flavonoids from red onion (*Allium cepa*), *Phytochemistry*. 47 (1998) 281–285.
- [38] H.J. Lee, H.J. Lee, E.O. Lee, S.G. Ko, H.S. Bae, C.H. Kim, K.S. Ahn, J. Lu, S.H. Kim, Mitochondria-cytochrome C-caspase-9 cascade mediates isorhamnetin-induced apoptosis, *Cancer Lett.* 270 (2008) 342–353.
- [39] E.H. Lee, H.J. Kim, Y.S. Song, C. Jin, K.T. Lee, J.S. Cho, Y.S. Lee, Constituents of the stems and fruits of *Opuntia ficus-indica* var. *saboten*, *Arch. Pharm. Res.* (12) (2003) 1018–1023.
- [40] M.H. Chavesa, N.F. Roque, M.C. Costa, Steroids and flavonoids of *Porcelia macrocarpa*, *J. Braz. Chem. Soc.* 15 (2004) 608–613.
- [41] K.D. Sen, D.W. Aksnes, G.W. Francis, NMR study of some Anthraquinones from rhubarb, *Magn. Reson. Chem.* 30 (1992) 359–363.
- [42] I. Tarascou, K. Barathieu, Y. Andre, I. Pianet, E. Dufourc, E. Fouquet, An improved synthesis of procyanidin dimers: regio and stereocontrol of the interflavan bond, *Eur. J. Org. Chem.* (23) (2006) 5367–5377.
- [43] F. Ahmad, M. Ali, P. Alam, New phytoconstituents from the stem bark of *Tinospora cordifolia* Miers, *Nat. Prod. Res.* 24 (2010) 926–934.
- [44] M.L. Flagg, G.A. Waechter, A.L. Davis, G. Montenegro, B.N. Timmermann, Two novel flavanones from *Greigia sphacelata*, *J. Nat. Prod.* 63 (2000) 1698–1691.
- [45] L.R. Chadwick, D. Nikolic, J.E. Burdette, C.R. Overk, J.L. Bolton, R.B. Breemen, R. Froehlich, H.H.S. Fong, N.R. Farnsworth, G.F. Pauli, Estrogens and congeners from *Spent Hops* (*Humulus lupulus*), *J. Nat. Prod.* 67 (2004) 2024–2032.
- [46] S.R. Ko, J. ChoK, K. Suzuki, Enzymatic preparation of ginsenosides Rg2, Rh1 and F1, *Chem. Pharm. Bull.* 51 (2003) 404–408.
- [47] T. Mosmann, Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *J. Immunol. Methods* 65 (1983) 55–63.
- [48] U. Schauer, I. Krolkowski, C.H. Rieger, Detection of activated lymphocyte subsets by fluorescence and MTT staining, *J. Immunol. Methods* 116 (1989) 221–227.
- [49] K. Ljung, L. Lioussanne, G. Sandberg, Developmental regulation of Indole-3-acetic acid turnover in scots pine seedlings¹, *Plant Physiol.* 125 (2001) 464–470.