

Two new species of *Paramphidelus* Andrassy, 1977 (Nematoda: Alaimidae) from Manipur, India

Mahlaqa CHOUDHARY and M. Shamim JAIRAJPURI *

Section of Nematology, Department of Zoology, Aligarh Muslim University, Aligarh-202001, India.

SUMMARY

Two new species of *Paramphidelus* from India are described. *Paramphidelus saccus* n. sp. is 0.8-0.9 mm long, has small tennis-racket shaped amphids at 18-20 μm from the anterior extremity and an anterior uterine sac. *Paramphidelus samii* n. sp. is 1.2-1.3 mm long, has funnel-shaped amphids at 10-12 μm from the anterior extremity, amphidelphic asymmetrical gonads and heavily sclerotized vagina.

RÉSUMÉ

Deux nouvelles espèces de *Paramphidelus* Andrassy, 1977 (Nematoda : Alaimidae) découvertes à Manipur, Inde

Deux nouvelles espèces de *Paramphidelus* découvertes en Inde sont décrites. *Paramphidelus saccus* n. sp. a une longueur de 0,8 à 0,9 mm et de petites amphides en forme de raquette situées à 18-20 μm de l'extrémité antérieure et un sac utérin antérieur. *Paramphidelus samii* n. sp. a une longueur de 1,2 à 1,3 mm, des amphides en forme d'entonnoir situées à 10-12 μm de l'extrémité antérieure ainsi que des gonades amphidelphiques asymétriques et un vagin fortement sclérotisé.

Soil samples collected from Manipur, India yielded two new species belonging to the genus *Paramphidelus* Andrassy, 1977 of the superfamily Alaimoidea, named *Paramphidelus saccus* n. sp. and *Paramphidelus samii* n. sp. Measurements were taken and observations made on specimens mounted in glycerine after being killed in hot 4% formalin and dehydrated by the slow method.

Paramphidelus saccus n. sp.

(Fig. 1)

DIMENSIONS

Female (paratypes ; n = 3) L = 0.8-0.9 mm ; a = 42-51 ; b = 3.8-4.0 ; c = 13-14 ; c' = 6-10 ; V = 41-42 ;

Holotype (female). L = 0.8 mm ; a = 43 ; b = 4.0 ; c = 12 ; c' = 6 ; V = 41.

DESCRIPTION

Body arcuate, tapering towards extremities. Cuticle finely striated presumably only in the inner layers. Lip region offset, 3-4 μm wide or 1/6-1/5 of midbody width. Amphids small, tennis-racket shaped, at 18-20 μm or 5-6 lip widths from anterior extremity, amphidial apertures 1.5-2.0 μm wide. Oesophagus 190-233 μm long, expanded part occupying 19-24% of its length. Dorsal and three ventrosublateral oesophageal gland nuclei present. Nerve ring 105-120 μm from anterior extremity. Cardia conoid, 4-5 μm long. Reproductive system opistho-monodelphic 165-203 μm long. Vulva transverse ; vagina 4-6 μm long. Genital tube reflexed, oocytes 14-18 arranged in a single row except at tip. Anterior uterine sac 28-32 μm or 1.3-1.5 vulval body width long. Rectum about one anal body width long.

* Present address : Commonwealth Institute of Parasitology, St. Albans, Herts, England.




Fig. 1. *Paramphidelus saccus* n. sp. A : Entire female ; B & C : Anterior region ;
D : Basal region of oesophagus ; E : Female genital tract ; F : Female tail.

Tail elongate-conoid, 75-98 μm or 6-10 anal body widths long.

Male: Not found.

TYPE HABITAT AND LOCALITY

Soil around roots of rose, *Rosa sinensis* L., Imphal Hill, Imphal, Manipur.

TYPE SPECIMENS

Collected in August 1977.

Holotype: on slide *Paramphidelus saccus* n. sp./1 deposited in the Nematode Collection of the Zoology Department, Aligarh Muslim University Aligarh.

Paratypes: on slide *Paramphidelus saccus* n. sp./2-3 deposited in the same Collection. One paratype female deposited at Muséum national d'Histoire naturelle, Laboratoire des Vers, Paris, France.

DIFFERENTIAL DIAGNOSIS

With Andrassy's (1977) key, *P. saccus* n. sp. keys with *P. candidus* (Siddiqi & Basir, 1965) Andrassy, 1977 from which it differs in having a wider body and longer oesophagus; in size and shape of tails, shorter anterior uterine sac and slightly more posterior vulva ($a = 57-62$; $b = 4.6-4.7$; $c = 4.3-4.4$; tail elongate-attenuated; anterior uterine sac 2.33 body widths long; $V = 39-40$ in *P. candidus*). From *P. sylvaticus* (Siddiqi & Basir, 1965) Andrassy, 1977, *P. lemani* (Stefanski, 1914) Andrassy, 1977 and *P. paramonovi* (Eliashvilli, 1971) Andrassy, 1977, which also have an anterior uterine sac, it differs as follows: From *P. sylvaticus* it differs in having a smaller and wider body, in shape of amphids and position of amphidial apertures, and in shorter tail ($L = 1.12-1.29$ mm; $a = 70-80$; amphids cup-shaped with kidney-shaped apertures at 2.5 lip-widths from anterior extremity; $c = 6.7-8.0$ in *P. sylvaticus*). From *P. lemani* it differs in having a shorter body, more posteriorly located amphidial apertures, shorter tail and more posteriorly located vulva ($L = 1.23-1.39$ mm; amphidial apertures situated at 3.0-3.5 lip-widths from anterior extremity; $c = 4.7-6.2$; $V = 35-38$ in *P. lemani*). From *P. paramonovi* it differs in having a shorter and wider body; more anteriorly located amphidial apertures, shorter tail and more anteriorly located vulva ($L = 1.14-1.20$ mm; $a = 104-110$; amphidial apertures situated at 3-4 lip-widths from anterior extremity; $c = 19-20$; $V = 47-49$ in *P. paramonovi*).

From *P. pseudobulbosus*, which also has amphids 5-6 head widths from the head-end, it differs in having a wider body and longer tail and in presence of an anterior uterine sac and more posteriorly located vulva ($a = 77-100$; $c = 22-28$; anterior uterine sac absent; $V = 49-53$ in *P. pseudobulbosus*).

Paramphidelus samii n. sp.

(Fig. 2)

DIMENSIONS

Female (paratypes; $n = 4$) $L = 1.2-1.3$ mm; $a = 65-68$; $b = 4.8-6.5$; $c = 12-13$; $c' = 8-10$; $V = 41-43$.

Holotype (female). $L = 1.3$ mm; $a = 67$; $b = 4.8$; $c = 12$; $c' = 8$; $V = 41$.

DESCRIPTION

Female: Body regularly tapering towards both extremities, more curved posteriorly. Cuticle finely striated presumably only in the inner layers. Lip region continuous with body, 5-6 μm wide or 1/4-1/3 of midbody width. Amphids elongate funnel-shaped, at 10-12 μm or about two lip-widths from anterior extremity, amphidial apertures 3-4 μm wide. Oesophagus 230-270 μm long, expanded part occupying 25-29% of its length. Dorsal and three pairs of ventrosublateral oesophageal gland nuclei present. Nerve ring at 105-112 μm from anterior extremity. Cardia conoid, 2-3 μm long. Reproductive system amphidelphic but asymmetrical, occupying 195-210 μm of body length; anterior gonad shorter and poorly developed as compared to posterior gonad. Vulva transverse; vagina inclined posteriad, heavily cuticularized, 10-11 μm or more than half of vulval body width long. Genital tube reflexed, anterior ovary with 3-4 small oocytes, posterior ovary with 15-20 large ones. Rectum about one anal body-width long. Tail filiform, 102-105 μm or 8-10 anal body-widths long.

Male: Not found.

TYPE HABITAT AND LOCALITY

Soil around roots of papaya (*Carica papaya* L.), Singjamei, Imphal, Manipur.




Fig. 2. *Paramphidelus samii* n. sp. A : Entire female ; B : Anterior region, dorsal view ; C : Anterior region, lateral view ; D : Basal region of oesophagus ; E : Female genital tract ; F : Female tail.

TYPE SPECIMENS

Collected in December 1980.

Holotype: on slide *Paramphidelus samii* n. sp./1 deposited in the Nematode Collection of Zoology Department, Aligarh Muslim University, Aligarh, India.

Paratypes: on slide *Paramphidelus samii* n. sp./2-4, deposited in the same Collection. One paratype female deposited at Muséum national d'Histoire naturelle, Laboratoire des Vers, Paris, France.

DIFFERENTIAL DIAGNOSIS

With Andrassy's (1977) key *P. samii* n. sp. keys with *P. uniformis* (Thorne, 1939) Andrassy, 1977 from which it differs in having a wider body, more anteriorly located vulva and a much larger tail ($a = 83$; $V = 46$; $c = 22.5$ in *P. uniformis*). From *P. sylphus* (Thorne, 1939) Andrassy, 1977 and *P. hyans* (Thorne, 1939) Andrassy, 1977, it differs by having a shorter body, more anteriorly located vulva, unequally developed sexual branches and a slightly longer tail ($L = 2.0$ mm; $V = 55$; sexual branches equally developed, $c = 10.6$ in *P. hyans*). From *P. sylphus* it differs in having a much shorter but wider body, in shape of lip region, shape of amphids and position of amphidial apertures, in having a more anteriorly located vulva, unequally developed sexual branches and a longer tail ($L = 2.4$ mm; $a = 111$; lip region conical, amphids elongate-conoid with their amphidial apertures at a distance of four lip-widths from anterior extremity, sexual branches equally developed, $V = 57$; $c = 18$ in *P. sylphus*).

Accepté pour publication le 17 mars 1982.

The new species has been named after the late Mr. Samiuddin Choudhary.

ACKNOWLEDGEMENTS

We thank the Head of the Department of Zoology, for providing necessary laboratory facilities. The first author thanks CSIR, New Delhi for financial assistance.

RÉFÉRENCES

- ANDRÁSSY, I. (1977). Die Gattungen *Amphidelu*-Thorne, 1939. *Paramphidelus* n. gen. und *Etams phidelus* n. gen. (Nematoda : Alaimidae) *Opusc. zool., Bpest.*, 14 : 3-43.
- ELIASHVILI, T. S. (1971). Two new soil-inhabiting nematode species (*Amphidelus paramonovi* n. sp. and *Tylenchorhynchus georgiensis* n. sp.) of Eastern Georgia USSR. *Bull. Acad. Sci. Georgian SSR*, 61 : 213-216.
- DE MAN, J. G. (1876). Onderzoekingen over vrij in de aarde levende Nematode. *Tijdschr. ned. dierk. Vereen.*, 2 : 78-196.
- SIDDIQI, M. R. & BASIR, M. A. (1965). *Amphidelus sylvaticus* n. sp. (Nematoda : Alaimina) from India, with a key to the species of *Amphidelus*. *Nematologica*, 11 : 343-348.
- STEFÁNSKI, W. (1914). *Recherches sur la faune des nematodes libres du Bassin du Lemán*. Thèse, Geneva. 72 p.
- THORNE, G. (1939). A monograph of the nematodes of the superfamily Dorylaimoidea. *Capita zool.*, 8 : 1-261.