

Kinetoplast DNA minicircles of phloem-restricted *Phytomonas* associated with wilt diseases of coconut and oil palms have a two-domain structure

Michel Dollet ^{*}, Nancy R. Sturm ¹, Jean-Charles Ahomadegbe ², David A. Campbell ¹

CIRAD, TA 30/G, Campus International de Baillarguet, 34398 Montpellier Cedex 5, France

Received 1 August 2001; accepted 15 September 2001

First published online 26 October 2001

Abstract

We report the cloning and sequencing of the first minicircle from a phloem-restricted, pathogenic *Phytomonas* sp. (Hart 1) isolated from a coconut palm with hartrot disease. The minicircle possessed a two-domain structure of two conserved regions, each containing three conserved sequence blocks (CSB). Based on the sequence around CSB 3 from Hart 1, PCR primers were designed to allow specific amplification of *Phytomonas* minicircles. This primer pair demonstrated specificity for at least six groups of plant trypanosomatids and did not amplify from insect trypanosomatids. The PCR results were consistent with a two-domain structure for other plant trypanosomatids. © 2001 Published by Elsevier Science B.V. on behalf of the Federation of European Microbiological Societies.

Keywords: Conserved sequence block; Hartrot; Marchitez sorpressiva; Polymerase chain reaction; Plant trypanosomatid

1. Introduction

Some phloem-restricted trypanosomatids are specifically associated with sudden wilts of coconut (hartrot) and oil palm (marchitez sorpressiva) endemic to Latin America [1]. These trypanosomatids are transmitted through pentatomid bugs from the genus *Lincus* [2]; however, the natural plant reservoir(s) of the trypanosomatids that kill these two introduced plants is unknown.

The taxonomy of trypanosomatids isolated from plants is complicated by the observation that several monoxenous genera of nebulous membership known as 'insect trypanosomatids', namely *Leptomonas*, *Herpetomonas*, and *Crithidia*, can multiply in fruit [3]. All kinetoplastids

found in plants, including fruit, have typically been placed together under the designation *Phytomonas*. Thus there is a need to distinguish between trypanosomatids from plants and insects.

Several markers have been used to explore the relative classification within the phytomonads. Isoenzyme, RAPD and 5S rDNA analyses have defined a minimum of eight groups, currently designated A–H [4–6]. Studies of the SL-RNA gene have defined differences within phloem-restricted parasites [7].

Mitochondrial minicircles contain the genes for guide RNAs involved in the editing of mitochondrial RNA [8]. They have been useful molecular markers for taxonomic, phylogenetic, and diagnostic studies of trypanosomatids [9]. Minicircle templates are small and abundant, and have been exploited to create sensitive diagnostic and molecular epidemiological tools using PCR [10]. The kDNA minicircles of plant trypanosomatids show a range of different sizes [11–13]; however, sequences of minicircles of plant trypanosomatids have been obtained only for fruit, specifically tomato, isolates [14,15]. In this paper, we report the first minicircle sequence from a pathogenic, phloem-restricted *Phytomonas* sp. and explore the use of the minicircles from *Phytomonas* spp. as informative molecular tools.

* Corresponding author. Tel.: +33 4 67 59 39 22;

Fax: +33 4 67 59 37 98.

E-mail address: dollet@cirad.fr (M. Dollet).

¹ Present address: Department of Microbiology, Immunology and Molecular Genetics, UCLA School of Medicine, University of California, 10833 Le Conte Avenue, Los Angeles, CA 90095-1747, USA.

² Present address: Unité des Marqueurs Génétiques des Cancers, Département de Biologie Clinique, Institute Gustave Roussy, 94805 Villejuif Cedex, France.

2. Materials and methods

2.1. Trypanosomatids

The isolate Hart 1 [11] was obtained from a coconut tree affected with hartrot in Saut Sabbat, French Guiana. The previously described *Phytomonas* spp. isolates used for PCR amplification are: Ps1G [16]; Lima.Pe, Berg.Br, E.hi.Sur, Mand.sc.Br, Tom.Sp, E.M.1, E.hi.Ind, E.hi.Sen, Mani.Br, E.hi.Vz, E.hy.Gu, Hart 1, Mar 1 and Ascl.cu.Vz [6]; Aca [11]; Mar 6 [7]; Blepharodon, Rhabdadenia, Trifolium and Amaranthus [5]. Previously undescribed Venezuelan isolates from *Euphorbia heterophila* (E.het.Vz) and *Euphorbia prostrata* (E.pro.Vz) were collected by M.D. from San Augustin, Estado Sucre in 1991 and Puerto Chama, Estado Zulia in 1992, respectively.

The following insect trypanosomatid DNAs were used as controls: *Tangerine* (São Paulo TCC 237), *Leptomonas mirabilis* (ATCC 30263), *Leptomonas seymouri* (ATCC 30220), *Crithidia oncopelti* and *Crithidia luciliae* (kindly provided by Theo Baltz, University of Bordeaux II), *Crithidia deanei* (ATCC 30255), *Crithidia acanthocephali* (ATCC 30251), *Herpetomonas muscarum* (Liverpool School of Tropical Medicine), *Herpetomonas megaseliae* (ATCC 30209) and *Herpetomonas samuelpessoai* (kindly provided by G. Riou, Institute Gustave Roussy).

2.2. kDNA extraction, cloning, sequencing and analysis

kDNA for cloning was obtained through fractionation in a CsCl-propidium iodide gradient as described previously [11]. Minicircles were cut with *Mbo*I (one cut), and cloned into the *Bam*HI site of the plasmid pBluescript SK+. Candidate clones were purified, analyzed by digestion with *Mbo*I, and screened for the presence of inserts corresponding to the size of Hart 1 minicircles [11,12]. A 0.8-kb *Pst*I fragment was subcloned to allow for complete sequencing of the minicircle clone. Template plasmids were purified using the QIAquick PCR Purification Kit (Qiagen) and sent to Davis Sequencing for dideoxy sequencing. Sequence information was compiled and analyzed using the UW GCG package.

2.3. PCR amplification

Synthetic oligonucleotides used for *Phytomonas* minicircle-specific amplifications from the CSB-3 motif were: mcCSB3a, 5'-GGTTT TTTAG GGGTT GGTAT AAT; mcCSB3b, 5'-TACCA ACCCC TAAAA AACCC C; mcCSB3aa, 5'-AAGGT TTTT AGGGG TTGGT ATA; mcCSB3bb, 5'-GGTAC CAACC CCTAA AAAAC CC. PCR reactions were carried out using reagents from the Qiagen PCR Core Kit. Amplification was obtained routinely from 10 ng of total cell DNA or directly from 1/1000 dilutions of guanidine isothiocyanate cell lysates (4×10^8 cells ml⁻¹) into H₂O. The cycle profile used was

94°C for 2 min; 30 cycles of 94°C for 15 s, 60°C for 15 s, 72°C for 1 min; 72°C for 9 min. Reaction products were separated on 1.0% SeaKem LE agarose gels in 1×TBE buffer.

3. Results and discussion

Sequence characterization of a cloned minicircle from the Hart 1 isolate of *Phytomonas* sp. revealed a molecule of 1604 bp (GenBank accession No. AF397906), whose features are summarized in Fig. 1A. Analysis of the sequence identified two conserved regions (CR) of approximately 220 bp, containing variants on the conserved sequence block (CSB) 1, 2 and 3 motifs found in all Trypanosomatidae [17]. The conserved regions extended upstream from CSB 3 and were nearly identical, with minor variability in the area 27–56 bp upstream from CSB 1 (data not shown). The spacing among the CSB elements in Hart 1 was shorter than described for tomato fruit isolates [14,15], and more closely approached the spacing found in *Trypanosoma brucei*.

The Hart 1 minicircle structure is similar to that described in *Phytomonas serpens* [15], with one variable region showing a strong bias in G and T residues and the other containing a potential guide RNA gene in the vicinity of the *Mbo*I site. Unlike *P. serpens*, the sequence between CSB 1 and CSB 3 is identical for both CRs in the Hart 1 minicircle.

Based on the alignment of the Hart 1 minicircle CSBs

Table 1
Plant trypanosomatid minicircle and PCR product sizes

Group [4–6]	Isolate	Size (kb)	PCR (kb)
A	1G	1.44 [15]	0.76
	Berg	ND ^a	0.7
	Lima	1.45 [11]	0.76
B	Aca	1.66 [11]	0.8
	E.hi.Sur	ND	0.86
	Blephar.	ND	0.74
	Rhabda.	ND	0.85
C	Mand.sc.Br	1.62 [11]	0.86
	Tom.Sp	1.64 [13]	0.9
	Trifolium	ND	0.85
	Amaran.	ND	0.84
D	E.hi.Ind	2.30 [11]	– ^b
	E.hi.Sen	2.20 [11]	–
	E.M.1	2.8 [11]	1.5
F	Mani.Br	ND	0.74
G	E.hy.Gu	1.30 [11]	0.68
	E.hi.Vz	ND	0.68
H-I	Hart 1	1.6 [11]	0.82
H-IIa	Mar 1	1.8 [13]	0.875
H-IIb	Mar 6	1.8 [13]	0.875
I	Ascl.cu.Vz	0.85 [11]	–
ND	E.pro.Vz	ND	0.64
	E.het.Vz	ND	0.85

^aND, not determined.

^b–, not amplified.




Fig. 2. Amplification of *Phytomonas* spp. minicircle domains. A: Strains of known minicircle size. B: Strains of unknown minicircle size and of insect trypanosomatids. Primers mcCSB3a and mcCSB3b were used for the reactions, and the products separated through a 1% agarose gel containing ethidium bromide. E.M.1 was included as a positive control. For the phytomonads, the 5S group (uppercase letters, with SL RNA subgroups [7]) and RAPD (lower case letters) designations are shown underneath. Lanes flanking each gel are molecular size markers shown in base pairs (1 kb Plus DNA Ladder, Gibco BRL).

yield products (Fig. 2B), as expected for some based on their minicircle sequence (Fig. 1B). Thus, the primer set mcCSB3a and mcCSB3b does not amplify minicircles from insect trypanosomatids, but yields products for most plant trypanosomatids. We will pursue the direct cloning of minicircles from the three refractory isolates to examine their sequences.

The Hart 1 minicircle sequence represents the first minicircle data from a pathogenic, phloem-restricted member of the *Phytomonas* genus. Our analyses of the nucleotides in the conserved region, and specifically in the region of CSB 3, have allowed the design of primer sets for amplification from many, but not all, of the phytomonad groups. Our future goals include the use of minicircles to distinguish between true plant trypanosomatids and opportunistic insect trypanosomatids, as well as among three types of plant trypanosomatids found in fruit, latex, or phloem. Further studies of minicircle amplification products will be useful for determining the relationships be-

tween close isolates using the sequence information contained in the CRs [18].

Acknowledgements

D.A.C. was a grantee of CIRAD. We thank Theo Baltz, Erney Camargo, Peter Kastelein, Guy Riou and Manuel Sanchez-Moreno for providing cell lines.

References

- [1] Dollet, M. (1984) Plant diseases caused by flagellate protozoa (*Phytomonas*). Annu. Rev. Phytopathol. 22, 115–132.
- [2] Desmier de Chenon, R. (1984) Recherches sur le genre *Linicus* Stal, Hemiptera Pentatomidae Discocephalinae, et son rôle éventuel dans la transmission de la marchitez du palmier à huile et du hartrot du cocotier. Oleagineux 39, 1–6.
- [3] Conchon, I., Campaner, M., Sbravate, C. and Camargo, E.P. (1989) Trypanosomatids, other than *Phytomonas* spp., isolated and cultured from fruit. J. Protozool. 36, 412–414.
- [4] Muller, E., Gargani, D., Banuls, A.L., Tibayrenc, M. and Dollet, M. (1997) Classification of plant trypanosomatids (*Phytomonas* spp.): parity between random-primer DNA typing and multilocus enzyme electrophoresis. Parasitology 115, 403–409.
- [5] Serrano, M.G., Camargo, E.P. and Teixeira, M.M.G. (1999) *Phytomonas*: analysis of polymorphism and genetic relatedness between isolates from plants and phytophagous insects from different geographic regions by RAPD fingerprints and synapomorphic markers. J. Eukaryotic Microbiol. 46, 618–625.
- [6] Dollet, M., Sturm, N.R., Sánchez-Moreno, M. and Campbell, D.A. (2000) 5S ribosomal RNA sequences define at least eight groups of plant trypanosomatids (*Phytomonas* spp.): phloem-restricted pathogens form a distinct section. J. Eukaryotic Microbiol. 47, 569–574.
- [7] Dollet, M., Sturm, N.R. and Campbell, D.A. (2001) The spliced leader RNA gene array in phloem-restricted plant trypanosomatids (*Phytomonas*) partitions into two major groupings: epidemiological implications. Parasitology 122, 289–297.
- [8] Sturm, N.R. and Simpson, L. (1990) Kinetoplast DNA minicircles encode guide RNAs for editing of cytochrome oxidase subunit III mRNA. Cell 61, 879–884.
- [9] Morel, C., Chiari, E., Camargo, E.P., Mattei, D.M., Romanha, A.J. and Simpson, L. (1980) Strains and clones of *Trypanosoma cruzi* can be characterized by pattern of restriction endonuclease products of kinetoplast DNA minicircles. Proc. Natl. Acad. Sci. USA 77, 6810–6814.
- [10] Sturm, N.R., Degrove, W., Morel, C. and Simpson, L. (1989) Sensitive detection and schizodeme classification of *Trypanosoma cruzi* cells by amplification of kinetoplast minicircle DNA sequences: use in diagnosis of Chagas' disease. Mol. Biochem. Parasitol. 33, 205–214.
- [11] Ahomadegbe, J.C., Dollet, M., Coulaud, D., Gargani, D. and Riou, G. (1990) Kinetoplast DNA permits characterization of pathogenic plant trypanosomes of economic importance. Biol. Cell 70, 167–176.
- [12] Ahomadegbe, J.C., Dollet, M. and Riou, G. (1992) Kinetoplast DNA from plant trypanosomatids responsible for the hartrot disease in coconut trees. Biol. Cell 74, 273–279.
- [13] Muller, E., Ahomadegbe, J.C., Coulaud, D., Gargani, D., Fernandez-Becerra, C. and Dollet, M. (1995) Variability of kinetoplast DNA from plant trypanosomatids responsible for hartrot and marchitez diseases. Phytopathology 85, 942–947.
- [14] Sa-Carvalho, D.S., Perez Morga, D. and Traub Cseko, Y.M. (1993) Characterization of kinetoplast DNA from *Phytomonas serpens*. J. Eukaryotic Microbiol. 40, 317–322.

- [15] Maslov, D.A., Hollar, L., Haghghat, P. and Nawathean, P. (1998) Demonstration of mRNA editing and localization of guide RNA genes in kinetoplast-mitochondria of the plant trypanosomatid *Phytomonas serpens*. *Mol. Biochem. Parasitol.* 93, 225–236.
- [16] Da Silva, J.B.T. and Roitmann, I. (1990) Growth of *Phytomonas serpens* in a defined medium; nutritional requirements. *J. Protozool.* 37, 521–523.
- [17] Ray, D.S. (1989) Conserved sequence blocks in kinetoplast minicircles from diverse species of trypanosomes. *Mol. Cell. Biol.* 9, 1365–1367.
- [18] Yurchenko, V., Kolesnikov, A.A. and Lukes, J. (2000) Phylogenetic analysis of Trypanosomatina (Protozoa: Kinetoplastida) based on minicircle conserved regions. *Folia Parasitol.* 47, 1–5.