

ANTHELMINTIC PROPERTY OF ARECANUT (*Areca catechu* Linn.) - A REVIEW

S. Keshava Bhat¹, S. Mythri² and D. Ashwin²

INTRODUCTION

Areca palm, *Areca catechu* Linn., belongs to the family Arecaceae is mainly distributed in South and South East Asia, especially in India, China, Bangladesh, Indonesia, Myanmar, Thailand, Malaysia, Vietnam, Philippines, etc. (Arjungi, 1976). It is an evergreen palm with an erect stem reaching more than 20 m height. The fruit is a drupe, round or ovoid in shape with 3 - 4 cm diameter when fully grown. There is a central ruminant endosperm or seed covered by a thin pericarp or husk. The pericarp is green in colour when immature and yellowish brown when ripe. The seed or nut is hard and light brownish in colour when mature. It has a characteristic astringent and slightly bitter in taste (Ananda, 2004).

The endosperm of areca fruit, which is commonly called as arecanut or betelnut or supari, is generally marketed after processing. In India, two types of processing, one called as 'red supari' and another as 'white supari' are in vogue. The former type is obtained by boiling and drying dehusked unripe arecanuts at different stages of maturity, whereas the latter type is obtained by drying ripe arecanuts and dehusking afterwards (Selvan *et al.*, 2004).

Among several diseases and parasitic infestations, helminth parasites cause

considerable problems for human beings and livestock including poultry. The helminth parasites which infest the intestine are Cestodes (tapeworms); Nematodes (hookworms and roundworms) and Trematodes (flatworms or flukes) (Mali and Mehta, 2008). Many modern synthetic medicines though very effective against these parasites, they also cause a number of side effects. Moreover, indiscriminate use and incorrect dosage of such drugs lead to the development of resistance to such compounds (Borgsteede, *et al.*, 1997; Gill, 1996; Yadav, 1990; Easwaran *et al.*, 2009).

Since ancient times, arecanut is being used for mastication in several countries as it is believed to have numerous medicinal and anthelmintic properties (Aman, 1969; Rao, 1982; Oxenham *et al.*, 2001; Jaiswal *et al.*, 2011; Peng *et al.*, 2015). World Health Organization (2009) has listed out as many as 25 different beneficial effects of *A. catechu* on mankind. Amudan *et al.* (2012) have reviewed the phytochemical and pharmacological potential of the seeds of *A. catechu* and suggested for further detailed studies to identify the biochemical compounds responsible for such actions. Tagboto and Townson (2001) have reviewed the antiparasitic properties of several medicinal plants and suggested to validate and work further on those aspects.

¹ARDF®, Varanashi Towers, Mission Street, Mangaluru: 575001, Karnataka.

²Kannur Dental College, Anjarakkandy: 670 612, Kannur Dt., Kerala.

Chemical constituents of arecanut

Shivashankar *et al.* (1969) reported that the nuts or the endosperm of arecanut contains: Polyphenols (including flavonoids and tannins) 11.1 - 29.8%; Polysaccharides 17.3 - 25.7%; Proteins 6.2 - 9.4%; Fats 8.1 - 15.1%; Fibres 8.2 - 15.4%; Alkaloids 0.11 - 0.24% and Minerals 1.1 - 2.5% (including calcium 0.05%, phosphorous 0.13% and iron 1.5 mg/100 g). Arecanut also contains Vitamin B6 (286.9 mg%) and Vitamin C (416.2 mg%). The alkaloid fraction of arecanut contains: arecoline, arecaidine, guvacoline and guvacine, the first one being the primary pharmacologically active component. Polyphenols decrease with maturity, whereas polysaccharides, fat, fibre and alkaloid contents increase with maturity of the nuts (Mathew *et al.*, 1964).

The Fatty acid composition of arecanut are: lauric acid (19.5%), myristic acid (46.2%), palmitic acid (12.7%), oleic acid (6.2%), linoleic acid (5.4%), hexadecenoic acid (7.2%) and minor proportions of stearic acid, decanoic acid and monoethylenic acids (Pathak and Mathur, 1954).

Anthelmintic properties of arecanut

Crude drugs, derived from plants, though less effective in curing the disease, are relatively safe and biodegradable compounds and cause very less harm to environment. Large number of medicinal plants (Mali and Mehta, 2008; Mohan *et al.*, 2015; Veerakumari, 2015) and their seeds (Goswami *et al.*, 2013) have been recorded to have anthelmintic properties. The endosperm of arecanut is known for its anthelmintic properties since time immemorial and is being practiced as folk medicine to expel

intestinal worms in China (Hashimoto *et al.*, 1988) and India (Aman, 1969; Badanaje, 2008; Murthy *et al.*, 2011).

Several researchers have reported that the condensed tannins of plant origin are having potent anthelmintic properties (Athanasiadou *et al.*, 2001; Haslam, 1996; Klongsiriwet *et al.*, 2015; Min and Hart, 2003; Williams *et al.*, 2014a; Williams *et al.*, 2014b). Williams *et al.* (2014a) also observed that the condensed tannins cause significant damage to the cuticle and digestive tissues of the larvae of helminth parasites. Klongsiriwet *et al.* (2015) reported that the anthelmintic property of condensed tannins could be significantly enhanced by the addition of certain flavonoid monomers. Apart from tannins, arecoline is also reported to have anthelmintic property (Farnsworth *et al.*, 1985; Fabricant and Farnsworth, 2001). Mohammed *et al.* (2014) synthesized imidazole analogs of arecoline and tested their efficacy *in vitro* against the earthworm, *Eudrilus eugeniae* and found that the synthesized analogs had very effective anthelmintic properties. Several studies have been conducted on the anthelmintic properties of arecanut which has got good amount of tannin and fair amount of arecoline in it.

Valenciano (1980) studied the anthelmintic property of arecanut against the helminth parasites of dogs and found that the decoction of arecanut was very effective against these parasites. Two to four month old puppies when administered with the decoction of arecanut at the rate of 2cc per kg body weight, provided fair control against the roundworm, *Toxocara canis* and the tapeworm, *Dipylidium caninum* infestations in them.

Saluthan and Billacura (2005) studied *in vitro* the anthelmintic property of different solvent extracts of air dried seeds of arecanut against the common earthworm, *E. eugeniae* and found that the Hexane Extract, which contained only tannins and polyphenols, was most effective than Ethyl acetate and aqueous extracts against this nematode with a mean time for paralysis of 4.8 minutes and mean time for death of 38.8 minutes. They also reported that the Ethyl acetate and aqueous extracts contained not only tannins and polyphenols, but also alkaloids and flavonoids.

As an alternative to commercial parasiticides, Cargill *et al.* (2008) evaluated three ethnomedical plant products such as fruits of papaya, dried nuts of arecanut and leaves of pineapple to find out a cheaper and effective means of controlling the infestations of two gastrointestinal nematodal parasites such as *Trichuris* and *Ascaris* in pigs in Indonesia. They found that the pigs fed with a diet supplemented with dried arecanut powder at the rate of 4g/10kg live weight of pig once a week for 4 consecutive weeks eliminated *Trichuris* and *Ascaris* completely. The growth rate in pigs fed with a supplement of arecanut in their feed was increased by 82 g/day whereas in the control group the growth rate was decreased by 27 g/day.

Rajapakse, *et al.* (2009) studied the anthelmintic activity of nearly 15 medicinal plants against gastrointestinal nematodes infesting goats in Sri Lanka. Among these plants they found that the crude extracts of *A. catechu* fruit kernel and *Adhatoda vasica* leaves significantly reduced the number of worm eggs present in faecal matter of naturally infested

goats. In the group treated with areca crude extract at a dose of 6 mg/kg body weight at 14 days intervals during the test period of six months there was a significant decrease in egg counts in faecal matter from the second month of the treatment itself, and body weight significantly increased from the fifth month onwards when compared with that of the control group. They also suggested that the arecanut extract could be successfully incorporated in the control programmes against gastrointestinal nematodes infesting goats.

Jeyathilakan *et al.* (2010) evaluated *in vitro* the anthelmintic properties of three plant extracts such as *A. catechu*, *Erythrina indica* and *Zingiber officinale* against the liverfluke (trematode) *Fasciola gigantica* which infests domestic livestock in India and other tropical countries and compared the results with that of the common anthelmintic/flukicide drug Oxyclozanide. They found that the arecanut extract had 100% lethal effect at a concentration of 1% and above. Further, arecanut extract was found even better than Oxyclozanide in the treatment against *F. gigantica*. At 1% concentration of arecanut extract, the flukes died after 2 min of exposure, whereas at the same concentration of Oxyclozanide the flukes died only after 4 min of exposure. Other two plant derivatives were found to be less effective.

Tangalin (2011) studied the anthelmintic property of mature arecanut to chickens, sheep and goat and compared the results with that of commercial dewormers. The study conducted on chicken revealed that adult roundworms, tapeworms and parasite eggs were expelled successfully with a dose of 1 g of processed arecanut/kg body weight of the animal. In goat

and sheep, a dose of 1.5 g processed arecanut/kg body weight of the animal showed good results against parasitic worms. The results further revealed that the arecanut expelled both eggs and adults of roundworms and tapeworms, whereas the commercial dewormers such as Albendazole and Valbazen expelled only the adults of roundworms.

Not only the nuts, but also the roots of arecanut are having anthelmintic properties. Baby and Raphael (2014) evaluated *in vitro* the anthelmintic property of the roots of arecanut against the Indian common earthworm, *Pheretima posthuma* and compared the results with that of the standard drug Albendazole. They found that the ethanolic extract of the roots of arecanut were found to be even more potent than that of the standard drug Albendazole against *P. posthuma*. At a concentration of 25 mg/ml the arecanut extract induced paralysis after 24 minutes of exposure, whereas with the same concentration of Albendazole the time for inducing paralysis was only after 32 minutes of exposure. With arecanut extract, the time for death was after 36 minutes of exposure but with Albendazole death was not observed during the experimental period of 4 hours.

The major constituents of the larvicidal principles in the endosperm of arecanut were identified to be a mixture of fatty acids such as lauric, myristic, palmitic and oleic acids (Kiuchi, *et al.*, 1987). In *in vitro* studies conducted by them they also found that the combination of fatty acids and tannins of arecanut had strong larvicidal activity against dog roundworm, *T. canis*. According to them the widespread habit of betel chewing may be effective in the prevention of parasitic diseases in man.

The anthelmintic action of arecanut extract was mainly due to its ability to inhibit the formation of glutathione S-transferase in the body of the parasite which lead to the accumulation of toxic metabolites leading to paralysis and death of the parasite (Dhanraj and Veerakumari, 2015). In their study conducted on the trematode (flake) parasite, *Cotylophoron cotylophorum* of sheep and goat, maximum level of such inhibition was observed at a concentration of 0.5 mg/ml after 8 hrs of exposure to the ethanol extract of arecanut.

Arecanut extract is also having ovicidal properties against nematode eggs. Such studies were done by Barbieri *et al.* (2014) on the eggs of the sheep gastrointestinal nematode, *Haemonchus contortus*. In *in vitro* trial conducted by them using the acetone extract of arecanut seed they found that the hatchability of the eggs was inhibited by 50 % at a dose of 7.45 mg/ml of the extract.

CONCLUSION

Plants are being used for the treatment of several diseases all over the world since the beginning of human civilization. In recent years, there has been growing interest in the pharmacological and therapeutic uses of medicinal plants because of their effectiveness, safety and easy availability. The therapeutic uses of arecanut as anthelmintic are reviewed in this paper. The literature shows that there is ample opportunity for scientists and pharmacologists for further studies in this field and isolate and synthesize the active principles actually responsible for anthelmintic properties from this plant which is abundantly grown in India and several other countries.

REFERENCES

- Aman, 1969. *Medicinal secrets of your food.* - *Areca nut*, Published by: Secretary, Indo-American Hospital, NR Mohalla, Mysore-7, India. pp700-702.
- Amudhan, M. S., Begum, V.H. and Hebbar, K.B. 2012. A review on Phytochemical and Pharmacological potential of *Areca catechu* L seed. *IJPSR*, **3**: 4151-4157.
- Ananda, K.S. 2004. Botany. In: *Arecanut*. (Eds: D. Balasimha and V. Rajagopal), Central Plantation Crops Research Institute, Kasaragod, Kerala, India. pp7-50.
- Arjungi, K.N. 1976. *Areca nut: a review.* *Arzneimittelforschung (Drug Res)*. **26**: 951-956.
- Athanasiadou, S., Kyriazakis, I., Jackson, F. and Coop, R.L. 2001. Direct anthelmintic effects of condensed tannins towards different gastrointestinal nematodes of sheep: *in vitro* and *in vivo* studies. *Veterinary Parasitol.*, **99**: 205 - 219.
- Baby, A.A. and Raphael, K.R. 2014. Potential antimicrobial, anthelmintic and antioxidant properties of *Areca catechu* L. root. *Int. J. Pharma. Pharmaceut. Sci.* **6**: 486 - 489.
- Badanaje, S.B. 2008. *Arecanut - medicinal and alternative uses.* Arecanut Research and Development Foundation®, Varanashi Towers, Mission Street, Mangaluru: 575 001, India. 104p.
- Barbieri, A.M.E., Fornazari, B.C., Canova, E.B., Moreira, E.L. and Katiki, L.M. 2014. Effectiveness of *Areca catechu* Linn against *Haemonchus contortus* *in vitro* egg hatch assay. *B. Industr. Anim., Nova Odessa*, **v.71**, (Suppl).
- Borgsteede, F. H. M., Peklder, J. J., Dercksen, D. P., Sol, J., Vellema, P., Gaasenbeek, C. P. H. and van der Linden, J. N. 1997. A survey of anthelmintic resistance in nematodes of sheep in the Netherlands. *Vet. Quart.* **19**: 167 - 172.
- Cargill, C., Syahputra, T., Damriyasa, M. 2008. *Feeding papaya fruits and betel nuts to reduce parasite burdens and increase growth rate in pigs.* Project Final Report, Australian Centre for International Agricultural Research (ACIAR), Canberra, Australia. 11p.
- Dhanraj, K.M. and Veerakumari, L. 2015. Effect of ethanol extract of *Areca catechu* and *Syzygium aromaticum* on glutathione-S-transferase of *Cotylophoron cotylophorum*. *World J. Pharma. Pharmaceut. Sci.*, **4**: 1117 - 1125.
- Easwaran, C., Harikrishnan, T.J. and Raman, M. 2009. Multiple anthelmintic resistance in gastrointestinal nematodes of sheep in Southern India. *Veterinarski Archiv*, **79**: 611 - 620.
- Farnsworth, N.R., Akerele, O., Bingel, A.S., Soejarto, D.D. and Guo, Z. 1985. Medicinal Plants in Therapy. *Bull. World Health Organ.*, **63**: 965 - 981.
- Fabricant, D.S. and Farnsworth, N.R. 2001. The value of plants used in Traditional Medicine for drug discovery. *Environmental Health Perspectives*, **109**: 69 - 75.

- Gill, B. S. 1996. Anthelmintic resistance in India. *Vet. Paracitol.*, **63**: 173 - 176.
- Goswami, S., Nishad, S., Rai, M., Madhesiya, S., Malviya, A., Pandey, P., Gautam, V. and Yadav, S. 2013. Plant seeds used for anthelmintic activity: a review. *Ind. J. Res. Pharma. Biotech.*, **1**: 533 - 536.
- Hashimoto, Y., Kawanishi, K and Moriyasu, M. 1988. Forensic Chemistry of Alkaloids by Chromatographic analysis. pp 1 - 77. In: *The Alkaloids - Chemistry and Pharmacology* Vol 32 (Ed: A. Brossi), Academic Press Inc., San Diego, California. 449p.
- Haslam, E. 1996. Natural polyphenols (vegetable tannins) as drugs: possible modes of action. *J. Nat. Prod.* **59**: 205 - 215.
- Jaiswal, P., Kumar, P., Singh, V.K. and Singh, D.K. 2011. *Areca catechu* L.: A valuable herbal medicine against different health problems. *Res. J. Med. Plants*, **5**: 145 - 152.
- Jeyathilakan, N., Murali, K., Anandaraj, A. and Basith, S.A. 2010. *In vitro* evaluation of anthelmintic property of herbal plants against *Fasciola gigantica*. *Indian J. Anim. Sci.* **80**: 1070 - 1074.
- Kiuchi, F., Miyashita, N., Tsuda, Y., Kondo, K. and Yoshimura, H. 1987. Studies on crude drugs effective on visceral larva migrans. I. identification of larvicidal principles in Betelnuts. *Chem. Pharm. Bull.*, **35**: 2880 - 2886.
- Klongsiriwet, C., Quijada, J., Williams, A.R., Harvey, I.M., Williamson, E.M. and Hoste, H. 2015. Synergistic inhibition of *Haemonchus contortus* exsheathment by flavonoid monomers and condensed tannins. *Int. J. Parasitol. Drugs Drug Resist.*, **5**: 127 - 134.
- Mali, R. G. and Mehta, A. A. 2008. A review on Anthelmintic Plants. *Natural Product Radiance*, **7**: 466 - 475.
- Mathew, A.G., Venkataramu, S.D. and Govindarajan, V.S. 1964. Studies on arecanut: part 1. Changes in chemical composition and physical characteristics of nuts with maturity. *Indian J Tech.*, **2**: 90 - 96.
- Min, B.R. and Hart, S.P. 2003. Tannins for suppression of internal parasites. *J. Anim. Sci.* **81** (E. Suppl.2): E102 - E109.
- Mohammed, M., Arul, K., Anjana, A.K. and Remya, K. 2014. Synthesis, docking studies and Pharmacological evaluation of imidazole analogues of arecoline. *Int. J. Curr. Pharmacut. Res.*, **6**: 22-26.
- Mohan, C., Saxena, N. and Fozdar, B.I. 2015. Activity of indigenously known angiospermic plants against common GI parasites of livestock. *World J. Pharma. Phamaceut. Sci.*, **4**: 1652 - 1667.
- Murthy, P.K., Joseph, S.K. and Murthy, P.S.R. 2011. Plant products in the treatment of filariasis and other helminth infections and assay systems for antifilarial/anthelmintic activity. *Planta Med.*, **77**: 647 - 661.
- Oxenham, M.F., Locher, C., Cuong, N.L. and Thuy, N.K. 2002. Identification of *Areca catechu* (Betel nut) residues on the dentition of Bronze age inhabitants of Nui

- Np, Northern Vietnam. *J. Archaeol. Sci.*, **29**: 1-7.
- Pathak, S.P. and Mathur, S.S. 1954. The component acids and glycerides of arecanut (*Areca catechu*) fat. *J. Sci. Food Agri.*, **5**: 461 - 465.
- Peng, W., Lie, Y.J., Wu, N., Sun, T., He, X.Y., Gao, Y.X. and Wu, C.J. 2015. *Areca catechu* (Arecaceae): A review of its traditional uses, botany, phytochemistry, pharmacology and toxicology. *J. Ethnopharma.*, **164**: 340-356.
- Rao, M.M. 1982. Introduction. In: *The Arecanut Palm*. (Eds: K.V.A. Bavappa, M.K. Nair and T.P. Kumar). Central Plantation Crops Research Institute, Kasaragod, Kerala, India. pp1-9.
- Rajapakse, R.P.V.J., Fernando, W.I.T. and Dharmaratne, H.R.W. 2009. Anthelmintic activity of medicinal plants extract against gastrointestinal nematodes in naturally infested goats. *International Symposium on sustainable improvement of animal production and health (Synopses)*. FAO/IAEA, Vienna, Australia. 8-11 June. pp346 - 347.
- Salutan, G.L.H. and Billacura, M.P. 2015. Phytochemical screening and evaluation of the toxicity, antimicrobial and anthelmintic properties of the different extracts from the air-dried seeds of *Areca catechu* Linn. In: *Annual PAASE Meeting and Symposium (APAMS)*, Feb 5-7, 2015. De La Salle Univ., Manila, Philippines. POH-3-3.
- Selvan, M.T., Sivaraman, K. and Manojkumar, K. 2004. Economics, marketing and development. In: *Arecanut*. (Eds: D. Balasimha and V. Rajagopal). Central Plantation Crops Research Institute, Kasaragod, Kerala, India. pp259-301.
- Shivashankar, S., Dhanaraj, S., Mathew, A.G., Murthy, S.S., Vyasamurthy, M.N. and Govindarajan, V.S. 1969. Physical and chemical characteristics of processed arecanuts. *J. Food Sci. Tech.*, **6**: 113-116.
- Tagboto, S. and Townson, S. 2001. Antiparasitic properties of medicinal plants and other naturally occurring products. *Advances Parasitol.*, **50**: 199 - 295.
- Tangalin, M.G.G. 2011. Anthelmintic effects of processed mature betel nut as dewormer to native chicken and small ruminants (sheep and goats). *Int. Peer Reviewed J.* **1**: 230 - 243.
- Valenciano, L.A. 1980. The anthelmintic activity of betel nut (*Areca catechu*) against *Toxocara canis*. *MSAC Res. J.*, **6/7**: 30-38.
- Veerakumari, L. 2015. Botanical Anthelmintics. *Asian J. Sci. Tech.*, **6**: 1881 - 1894.
- WHO, 2009. *Areca catechu* L. pp 30 - 31. In: *Medicinal Plants of Papua New Guinea*, World Health Organization, Geneva, Switzerland, 302p.
- Williams, A.R., Fryganas, C., Ramsay, A., Muller-Harvey, I. and Thamsborg, S.M. 2014a. Direct anthelmintic effects of condensed tannins from diverse plant sources against *Ascaris suum*. *PLoS ONE* **9**(6): e99738.
- Williams, A.R., Ropiak, H.M., Fryganas, C.,

Desrués, O., Muller-Harvey, I. and Thamsborg, S.M. 2014b. Assessment of the anthelmintic activity of medicinal plant extract and purified condensed tannins against free-living and parasitic stages of

Oesophagostomum dentatum. *Parasites & Vectors*, 7: 518.

Yadav, C. L. 1990. Fenbendazole resistance in *Haemonchus contortus* of sheep. *Veterinary Records*, 126: 586.

**

**THE CASHEW NUT !
A WONDER NUT OF NATURE
HEALTHY YOU ARE WHEN YOU CONSUME IT.
WEALTHY YOU ARE WHEN YOU GROW IT.**

For more details - Read our publications :

1. The Cashew
2. Cashew Cultivation Science & Techniques
3. Cashew Varietal Wealth of India
4. Indian Cashew Industry

For further details Contact :

**THE DIRECTOR
DIRECTORATE OF CASHEWNUT & COCOA DEVELOPMENT
KERA BHAVAN, COCHIN - 682 011
KERALA**

Web: <http://dccd.gov.in>, E mail : dccd@nic.in

Phone : 0484-2377151 Fax: 0484-2377239