

Population Growth Potential of *Bracon brevicornis* Wesmael (Braconidae: Hymenoptera): A Life Table Analysis

T. SRINIVASAN^{1*} and C. MOHAN²

¹*Coconut Research Station, Tamil Nadu Agricultural University, Aliyarnagar, Tamil Nadu – 642 101, India

²ICAR-Central Plantation Crops Research Institute Regional Station, Kayankulam, Kerala – 690 533, India

(Received: 11 November 2016; accepted: 2 January 2017)

A laboratory experiment was conducted at the Biocontrol laboratory, Coconut Research Station, Aliyarnagar to study the life table parameters of *Bracon brevicornis* Wesmael, a parasitoid of the coconut black-headed caterpillar, *Opisina arenosella* Walker on its established laboratory host, *Corcyra cephalonica* Stainton. Studies at 28.3 ± 0.1 °C and $59.6 \pm 0.6\%$ r.h. revealed that the net reproductive rate (R_0) of *B. brevicornis* was 39.52 females/ female when reared on *C. cephalonica* larvae. The precise generation time (T) was 13.33 days. The intrinsic rate of natural increase (r_m) was 0.2758 which was slightly higher than the innate capacity for increase ($r_c = 0.2504$). The weekly multiplication rate was 6.893 numbers while 1561.83 females could be expected in the F_2 generation. Higher net reproductive rate, coupled with shorter population doubling time of *B. brevicornis* indicate the efficacy of the parasitoid as a suitable candidate for the management of coconut black headed caterpillar, *O. arenosella*, its intended host under field conditions.

Keywords: population growth potential, life table, *Bracon brevicornis*.

Biological control agents are gaining increased attention in the recent years due to their environmental feasibility, eco-friendliness and sustainability. Coconut, *Cocos nucifera* L. is a perennial plantation cum oil seed crop grown in India with more than 10 million people depending upon it for livelihood (Nampoothiri and Thomas, 2000). Coconut is grown in an area of 1.97 million ha with an annual production of 20,439 million tonnes (CDB, 2016). Among the several pests attacking the palm, the black headed caterpillar, *Opisina arenosella* Walker is one of the serious lepidopterous pests causing damage by scrapping the photosynthetic material from the abaxial surface of the leaves and making galleries of excreta on the under surface of the leaves. This leads to reduction in the photosynthetic efficiency and thereby cause yield reduction to the tune of 30-40 per cent (Mohamed, 1980). Being a tree crop and attaining a height of even 25 to 40 feet, the black headed caterpillar management poses a serious hurdle in that, the plant protection measures such as insecticidal sprays do more harm than good to the plant protection worker as well as to the ecosystem besides being harmful to non-target organisms and natural enemies. Species belonging to the Order Hymenoptera have been recorded mainly as par-

* Corresponding author; e-mail: entosrini@gmail.com

asitoids of *O. arenosella*. Thus biocontrol agents, especially the larval parasitoids *Bracon brevicornis* Wesmael and *Goniozus nephantidis* Muesebeck serve as effective parasitoids for the management of *O. arenosella* under field conditions (Cock and Perera, 1987). Adult females of *B. brevicornis* lay about 25 eggs on the larva of *O. arenosella*. The eggs hatch out in 2-3 days and the emerging larvae survive on the host larvae (*O. arenosella*) for 2-3 days. The pupal stage lasts about 3-4 days and the emerging adults go in search of new host larvae (Narendran, et. al., 1980; Rukshana and Majeed, 2014). Release of larval parasitoids alone can bring about 83 per cent reduction in the pest population (Nair, 1993). This paper is an attempt to study the life table parameters of the parasitoid, *B. brevicornis* on one of its laboratory host, *Corcyra cephalonica* Stainton under laboratory conditions.

Life table studies of parasitoids and predators provide necessary information on their effectiveness against target pests, rate of population build up, etc. under laboratory conditions which can be correlated to natural conditions (Muthukrishnan et al., 1995) and to schedule period of intervals for parasitoid releases. Life table computation helps in studying the population dynamics of the insect species and is an analytical tool which provide a description of the survivorship, development, etc, and reveal the maximum growth potential of a population (Gabre et al., 2004).

Materials and Methods

Rearing of C. cephalonica

To maintain the *C. cephalonica* culture, eggs were placed in coarsely broken sorghum (*Sorghum bicolor* L. Moench.) grains in plastic trays (39 cm dia. × 14 cm height). The setup was left undisturbed at room temperature (28.3 ± 0.1 °C and $59.6 \pm 0.6\%$ r.h) for about 45 to 55 days and the emerging adults were collected, paired in a open transparent glass beaker (10 cm dia × 23 cm height) for about two to three days. The female adults loosely lay their eggs inside the beaker which can be sieved through a 30 mesh sieve. The collected eggs are placed in fresh broken sorghum grains for continuous rearing.

Rearing of Bracon brevicornis

Fully grown larva (from the above set up) forms the substratum for the braconid, *B. brevicornis*. Know number of full grown *C. cephalonica* larva were collected and placed on a tissue paper covering an open glass beaker (10 cm dia × 23 cm height). Immediately, a ghada cloth was placed on the larvae to arrest their movement to facilitate parasitisation by the parasitoids which were previously placed in the container. This is referred to as the Sandwich method (tissue paper – larvae – ghada cloth) that is widely followed for rearing *B. brevicornis* parasitoids (Jhansi, 1984). The tissue paper gives way for parasitisation of the larva placed above it and within 24 to 48 hrs, the larvae gets parasitized. After about 6 to 8 days, the ghada cloth was removed and the tissue paper containing the pupal stages of the parasitoid were placed in separate beakers (10 cm dia × 23 cm height) along with a cotton swab containing honey which serves as food for the emerging parasitoids.

Life table studies

In order to perform life table studies, the *C. cephalonica* larvae and *B. brevicornis* adults from the above culture were used. The laboratory studies were performed at room temperature 28.3 ± 0.1 °C and $59.6 \pm 0.6\%$ r.h. One pair of freshly emerged *B. brevicornis* was placed in a transparent glass vial (2 cm dia; 3.5 cm height) containing full grown 5th instar larva of *C. cephalonica* and plugged with cotton wool. A small cotton swab soaked in honey was placed inside the vial to serve as food for the parasitoids. The parasitoids were transferred to a new vial containing a full grown larva of *C. cephalonica* each day at a specific time until the female dies. Twenty five such vials (representing 25 replicates) were maintained and the experiment was continued until the females in all the 25 vials died naturally. The larval mortality, pupal mortality, sex ratio, longevity of male and female, etc. were studied along with other life table parameters viz., net reproductive rate (R_o), approximate generation time (T_c), intrinsic rate of natural increase (r_m), innate capacity for increase (r_c), finite rate of increase (λ), mean generation time (T), population doubling time (t), weekly multiplication rate (WMR) and hypothetical female in F_2 generation (R_o^2) as per the methodology of Venkatesan et al. (2004). The formulae for estimating the life table parameters are listed below (Table 1). The age specific survival (l_x) and age specific fecundity (m_x) at each pivotal age (x) was worked out daily for the entire reproductive period to prepare fertility table.

Results

Development of immature stages of *B. brevicornis* on *C. cephalonica* from oviposition to adult emergence took 8 days at 28.3 ± 0.1 °C and $59.6 \pm 0.6\%$ r.h. The larval and pupal mortality was 6.08% and 10.25 %, respectively (Table 2). The sex ratio in the present study revealed a slightly male biased population (0.9). The female adults had a

Table 1

Formulae for assessment of life table parameters

Life table parameters	Denoted by	Formulae
Net reproductive rate	R_o	$\sum l_x m_x$
Approximate generation time	T_c	$\sum x \times l_x m_x / \sum l_x m_x$
Innate capacity for increase	r_c	$\log_e R_o / T_c$
Intrinsic rate of natural increase	r_m	$e^{-r_m} \times \sum l_x m_x = 1$
Mean generation time	T	$\log_e R_o / r_m$
Finite rate of increase	λ	antilog e^{r_m}
Weekly multiplication rate	WMR	λ^7
Hypothetical female in F_2 generation	R_o^2	R_o^2
Population doubling time	DT	$\log_2 2 / r_m$

Table 2Biological parameters of *B. brevicornis* on *C. cephalonica*

Parameters	Range	Values
Larval mortality (%)	0.0–44.4	10.3 ± 2.2
Pupal mortality (%)	0.0–38.5	11.0 ± 2.1
Sex ratio (Female: Male)	0.1–6.3	1.2 ± 0.2
Female longevity (days)	6–31	20.4 ± 1.4
Male longevity (days)	1–7	3.2 ± 0.3

** Values represent mean ± standard error

longevity of 20.4 days while the males a meagre, 3.2 days. The life table and age specific fecundity was worked out (Table 3) based on which, the net reproductive rate (R_0) was found to be 39.52 females per female during its lifetime and the approximate generation time (T_c) was 14.68 days while, the precise generation time (T) stood at 13.33 days (Table 4). Thus the parasitoid has the capacity to multiply 39.52 times in a generation time of 13.33 days. The intrinsic rate of increase (r_m) was found to be slightly higher (0.2758) than the innate capacity for increase ($r_c = 0.2504$). The finite rate of increase (λ) was 1.318 nos. per day while, the weekly multiplication rate stood at 6.893 nos. and 1561.83 females could be expected in the F_2 generation.

Discussion

The development of immature stages of *B. brevicornis* on *C. cephalonica* from oviposition to adult emergence took 8 days at 28.3 ± 0.1 °C and $59.6 \pm 0.6\%$ r.h. and the sex ratio in the present study revealed a slightly male biased population (0.9). Male biased development was earlier reported by Singh et al. (2014) at 30 degree C. The female adults had a longevity of 20.4 days while the males a longevity of 3.2 days. Thus, augmentative parasitoid releases can be scheduled after 21 days after first release of parasitoids. Almost similar was the female longevity of *Bracon hebetor* (reared on *C. cephalonica*) (19.11 days), in experiments conducted by Farag et al. (2015), while the males had a longevity of 9.2 days. The net reproductive rate (R_0) in the present studies was found to be 39.52 females per female during its lifetime with an approximate generation time (T_c) of 14.68 days and a precise generation time (T) of 13.33 days. Thus the parasitoid has the capacity to multiply 39.52 times in a generation time of 13.33 days. Slightly higher intrinsic rate of increase (r_m) (0.2758) than the innate capacity for increase ($r_c = 0.2504$) suggest that the parasitoids have overlapping generations as opined by Southwood, 1976. Laboratory experiments by Farag et al. (2015) revealed differences in the net reproductive rate (R_0), intrinsic rate of increase (r_m), finite rate of increase (λ) and mean generation time (T) which were 30.6 female/ female/ life time, 0.19, 1.2133 and 18.09 days, respectively. Almost similar values for innate capacity for increase ($r_c = 0.2613$), intrinsic rate of increase ($r_m = 0.275$), finite rate of increase ($\lambda = 1.3168$), weekly multiplication rate (WMR = 6.86)

Table 3Life table and age specific fecundity of *B. brevicornis* on *C. cephalonica*

Pivotal age in days (x)	Proportion of survival of female at different stages (l_x)	No. of females produced per female at different ages (m_x)	$l_x m_x$	$x l_x m_x$
Immature stages 8 days				
9	0.92	2.04	12.7	16.9
10	0.92	3.43	11.7	31.6
11	0.88	4.45	11.2	43.1
12	0.88	5.86	10.2	61.9
13	0.80	5.20	10.1	54.1
14	0.80	7.80	9.1	87.4
15	0.72	2.56	9.1	27.6
16	0.72	2.22	8.1	25.6
17	0.68	2.53	7.5	29.2
18	0.64	3.13	7.0	36.0
19	0.56	0.93	6.9	9.9
20	0.52	3.77	6.4	39.2
21	0.52	5.62	5.4	61.3
22	0.52	1.46	4.4	16.7
23	0.40	1.50	4.6	13.8
24	0.36	3.00	4.0	25.9
25	0.36	3.78	3.0	34.0
26	0.36	2.00	2.0	18.7
27	0.16	6.25	2.9	27.0
28	0.16	4.00	1.9	17.9
29	0.08	2.50	2.3	5.8
30	0.08	1.50	1.3	3.6
31	0.04	3.00	1.5	3.7
32	0.04	3.00	0.5	3.8
			$\Sigma = 39.52$	$\Sigma = 580.28$

and population doubling time (DT = 2.52 days) were observed by Singh et al. (2014) in their life table experiments with *B. hebetor* at 30 degree C. However they have reported a higher net reproductive rate ($R_0 = 92.0$).

The higher female output and higher female longevity (up to 32 days) in the present study reveals that *C. cephalonica* possess the requisite nutritional quality for the growth and development of the larval parasitoid, *B. brevicornis* (Chandrababu et al., 1999). Earlier studies by Jhansi and Babu (2003) also revealed, *C. cephalonica* as a suitable host for mass rearing of the larval parasitoid, *Bracon brevicornis* with respect to oviposi-

Table 4Life table statistics of *B. brevicornis* on *C. cephalonica*

Parameters	Denoted by	Statistics
Net reproductive rate	R_o	39.52 females / female/ life time
Approximate generation time	T_c	14.68 days
Precise generation time	T	13.33 days
Innate capacity for increase	r_c	0.2504
Intrinsic rate of natural increase	r_m	0.2758
Finite rate of increase	λ	1.318 no./ day
Weekly multiplication rate	WMR	6.893 nos.
Hypothetical female in F_2 generation	(R_o^2)	1561.83
Population doubling time	DT	2.513 days

tion, hatchability, adult emergence and growth index. Thus, higher net reproductive rate, shorter population doubling time, higher female longevity, etc. indicate that *B. brevicornis* is an effective parasitoid for the management of black headed caterpillar in coconut ecosystems.

Acknowledgements

The financial assistance of the All India Coordinated Research Project (Palms), ICAR-Central Plantation Crops Research Institute, Kasaragod, Kerala, India is acknowledged.

Literature

- CDB (2016): (<http://www.coconutboard.gov.in/stat.htm>), Last modified: October, 15, 2016.
- Chandrababu, A., Gautam, R. D. and Garg, A. K. (1999): Biology of the ladybird beetle on aphid and mealybugs. *Ann. Pl. Prot. Sci.* 7, 13–18.
- Cock, M. J. W. and Perera, P. A. C. R. (1987): Biological control of *Opisina arenosella* Walker (Lepidoptera: Oecophoridae). *Biocontr. News Inf.* 8, 283–310.
- Farag, N. A., Ismail, I. A., Elbeherly, H. H. A., Abdel-Rahman, R. S. and Abdel Raheem, M. A. (2015): Life table of *Bracon hebetor* Say (Hymenoptera: Braconidae) reared on different hosts. *Intl. J. ChemTech. Res.* 8, 123–130.
- Gabre, R. M., Adham, F. K. and Chi, H. (2004): Life table of *Chrysomya megacephala* (Fabricius) (Diptera: Calliphoridae). *Acta Oecol.* 27, 179–183.
- Jhansi, K. and Babu, P. C. S. (2003): Comparative biology of *Bracon hebetor* Say in two host insects. *J. Appl. Zool. Res.* 14, 165–168.
- Jhansi, K. (1984): Studies on the *Bracon hebetor* (Say), a larval parasitoid of *Maruca testulalis* (Geyer). (Unpublished Master's thesis), Tamil Nadu Agricultural University, Coimbatore, India.
- Mohamed, U. V. K. (1980): Studies on the entomophagous insects associated with *Nephantis serinopa* Meyr. (A report of the work done under P.L. 480 research project). University of Calicut, Kerala.

- Muthukrishnan, J., Srinivasaperumal, S. and Selvan, M. S. (1995): Bioenergetics and life tables of selected parasitoids and predators. In: T. N. Ananthakrishnan (ed.): Biological Control of Social Forest and Plantation Crops Insects. Science Publishers: Lebanon. pp. 193–206.
- Nair, M. K. (1993): *Opisina arenosella* Wlk. The leaf eating caterpillar of coconut palm. Technical Bulletin No. 27, CPCRI, Kasaragod, India, 12 p.
- Nampoothiri, K. and Thomas, V. (2000): The present scenario of coconut, arecanut and cocoa in India. In: Proc. International Conference on Plantation Crops, PLACROSYM XIV, Hyderabad, India, 12-15 December, Souvenir, 8015 p.
- Narendran, T. C., Mohammed, U. V. K., Abdurahiman, U. C. and Kumar, V. V. S. (1980): Oviposition behaviour of *Bracon brevicornis* Wesmael (Hymenoptera: Braconidae). Agric. Res. J. Kerala, 18, 239–241.
- Rukshana, K. and Majeed, P. P. (2014): The effectiveness of biocontrol of *Opisina arenosella* (Lepidoptera: Oecophoridae) using *Bracon brevicornis* (Hymenoptera: Braconidae). J. Zool. Stud. 1, 1–3.
- Singh, D., Mathew, I. L. and Singh, R. P. (2014): Influence of different diets on intrinsic rate of natural increase (r_m) of *Bracon hebetor* (Say) (Hymenoptera: Braconidae) reared on *Corcyra cephalonica* Stainton (Lepidoptera: Pyralidae). Biolife, 3, 653–656.
- Southwood, T. R. E. (1976): Ecological Methods with Particular Reference to the Study of Insect Populations. Chapman and Hall, London, pp. 1–391.
- Venkatesan, A., Jalali, S. K., Murthy, K. S., Rabindra, R. J. and Rao, N. S. (2004): Comparative life table studies of *Goniozus nephantidis* (Muesback) (Hymenoptera: Bethyridae) on three different lepidopteran insect pests. Ann. Pl. Prot. Sci. 12, 5–8.