

2753
**Effect of *Paecilomyces lilacinus* for the control of burrowing nematode,
Radopholus similis on arecanut seedlings**

S. SUDHA, P. SUNDARARAJU* and ROHINI IYER

Central Plantation Crops Research Institute, Kasaragod - 671 124, Kerala

The fungus, *Paecilomyces lilacinus*, is a potential biocontrol agent for the control of plant parasitic nematodes. Sosamma *et al* (1990) studied the effect of *P. lilacinus* as a biocontrol agent

against burrowing nematode on betelvine. In the present investigation, an attempt has been made to study its effect against burrowing nematode on arecanut seedlings.

A pot culture experiment was initiated in arecanut seedlings variety Mangala, under green house conditions to evaluate the effect, if any, on burrowing nematode by *P. lilacinus*.

One year old arecanut seedlings, raised in sterilized soil, were transplanted in 45 x 30 cm earthen pots during January, 1991. Each pot contained 3.5 kg of methyl bromide fumigated sandy loam soil and the plants were maintained under green house conditions. There were 7 treatments, each replicated 5 times. The treatments were ;

- T1 - Control;
- T2 - Nematode alone;
- T3 - Fungus alone;
- T4 - Nematode first followed by fungus;
- T5 - Fungus first followed by nematode
- T6 - Nematode and fungus simultaneously

The experiment was laid out in a completely randomised block design and the treatments were imposed after 3 months. The nematode population, multiplied axenically on carrot discs, was used as inoculum source. In all the treatments the nematode inoculum(1000/plant) consisted of larval stages and adult females.

P. lilacinus was grown on wheat grains. For this, 250 g wheat grain was boiled in 250 ml of water, cooled and decanted. The cooked grains were distributed into five 250 ml flasks (50g) autoclaved at 15 lbs for 15 minutes. Upon cooling, the flasks were inoculated with 5 mm discs of PDA grown fungal mycelium (3 days old). Flasks were inoculated at room temperature (30°C) till the grains were completely covered with the fungal growth (4 weeks).

The fungal culture in each flask was divided into 5 parts and mixed with soil in each pot by gently raking and incorporating the fungal growth into it alongwith wheat grains. Nematode inoculum

Table 1. Effect of *Paecilomyces lilacinus* against *Radopholus similis* infesting arecanut seedlings.

(Mean of five replications)

Sl. No.	Treatment	Nematodes/10g root*	Nematodes/250g soil*	Root-lesion index*
1.	Nematode alone	5.1 (23.8)	23.2 (546.0)	3.4
2.	Nematode - fungus	2.5 (6.6)	13.8 (193.0)	2.8
3.	Fungus - nematode	1.0 (1.8)	11.5 (132.2)	1.2
4.	Nematode + fungus simultaneously	3.8 (14.8)	13.8 (193.0)	1.6
5.	Fungus alone	0.0	0.0	1.0
6.	Control	0.0	0.0	1.0
	C.D. (P=00.05)	1.01	2.57	0.85

* Transformation used is $\frac{1}{x}$ where x is the nematode count;
 Figures in parenthesis are the original values.

** 1 = No lesion; 1 to 2 = Moderate lesions ; 2 to 3 Severe lesions; 3 to 4 = Very severe lesions

was also mixed into the pot by gently raking the soil and mixing the nematode suspension into it. The experimental pots were maintained in green house at a temperature range of 27-30°C. Eight months after inoculation, plants were depotted carefully with intact root system and the plant growth parameters alongwith root lesion index were recorded. Both soil and root samples were collected from each treatment. Root samples were processed for assessing the nematode population according to procedures described by Koshy *et al.* (1975). Soil samples were processed by wet sieving and decanting method followed by modified Baermann funnel method.

The results showed that the inoculation of fungus first followed by nematode significantly reduced *R. similis* population in root compared to other treatments. The same trend was discernible in soil population also. The nematode population was

significantly low in treatments inoculated first with the fungus indicating that the fungus affected adversely nematode multiplication as well as root-lesion index. The present findings are in accordance with that of Sosamma *et al.* (1990) on betelvine.

ACKNOWLEDGEMENTS

The authors are highly thankful to the Director, CPCRI for providing facilities and Mr. K. Vijayakumar, Scientist, Senior Scale (Statistics) for the statistical analysis of the data.

REFERENCES

- Koshy, P.K., Sosamma, V.K., & Nair, C.P.R. (1975).
Indian J. Nematol. 5 : 26-35.
- Sosamma, V. K., Geetha, S.M. & Koshy, P.K. (1990).
Paper presented at the seminar on Bioagents in nematode management at IARI, New Delhi, July 13, 1990.