

The effect of flow rate at different pressures and temperatures on cocoa butter extracted from cocoa nib using supercritical carbon dioxide

E. K. Asep^{1,2} · S. Jinap^{1,2} · A. R. Russly¹ · M. H. A. Jahurul³ · Kashif Ghafoor⁴ · I. S. M. Zaidul⁵

Revised: 28 July 2015 / Accepted: 17 August 2015 / Published online: 24 May 2016
© Association of Food Scientists & Technologists (India) 2016

Abstract The effects of flow rate, different pressures and temperatures on cocoa butter extracted from cocoa nib using supercritical carbon dioxide (scCO₂) were investigated. The yield was analyzed for total fat content, triacylglycerol (TG) profile, and fatty acid (FA) profile. Extractions were carried out at pressures of 20 and 35 MPa, temperatures of 50 and 60 °C, and CO₂ flow rates of 0.5, 1, 2, 4 mL min⁻¹. The result shows that the yield of cocoa butter extract increased with increasing pressure, temperature, and flow rate and the optimum conditions for the maximum cocoa butter extraction were 35 MPa, 60 °C and 2 mL min⁻¹, respectively. TGs and FAs were found to be similar in composition to those of cocoa

butter obtained by conventional methods. The lower molecular weight TGs and FAs showed higher selectivity compared to higher molecular weight TGs and FAs.

Keywords Cocoa butter · scCO₂ extraction · Pressure · Temperature · Flow rate · Selectivity

Highlights

- Cocoa butter extraction from cocoa nib using supercritical carbon dioxide
- Effects of CO₂ flow rate, pressure, and temperature on cocoa butter extraction
- Cocoa butter quality base on extraction method
- Triglycerides in term of fatty acids constituents.

✉ S. Jinap
jinap@food.upm.edu.my; sjinap@gmail.com

I. S. M. Zaidul
zaidul@iiu.edu.my

¹ Food Safety Research Centre (FOSREC), Faculty of Food Science and Technology, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

² Institute of Tropical Agriculture, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

³ Faculty of Food Science and Nutrition, Universiti Malaysia Sabah, 88400 Kota Kinabalu, Sabah, Malaysia

⁴ Department of Food Science and Nutrition, King Saud University, Riyadh 11451, Saudi Arabia

⁵ Department of Pharmaceutical Technology, Faculty of Pharmacy, International Islamic University Malaysia, Kuantan Campus, 25200 Kuantan, Pahang, Malaysia

Introduction

Cocoa butter is unique among vegetable fats due to its composition and crystallization behavior. Cocoa bean contains substantial amounts (55 %) of cocoa butter (Saldaña et al. 2002; Rodríguez et al. 2014). There is a high demand for quality cocoa butter, especially for use in foods, nutraceuticals, cosmetics and pharmaceuticals (Asep et al. 2013; Jahurul et al. 2014, 2015). In comparison with the conventional method, supercritical fluid extraction (SFE) is a feasible and a potential method to be implemented in the cocoa industry to produce higher yield and better quality cocoa butter due to its rapid, non-toxic, environmental friendly, contamination-free and easily manipulated conditions (Jahurul et al. 2012; Jinap et al. 2013).

In recent years, there has been increasing interest in the development of supercritical fluid technology and its application. Recently the uses of SFE are widely increasing for the extraction, purification, crystallization, fractionation, and nanoparticle formation in food, nutraceutical and pharmaceutical industries (McHugh and Krukons 1994; Reverchon and Adami 2006; Campardelli et al. 2012; Jahurul et al. 2012; Knez et al. 2014). Compared to traditional organic solvents, supercritical fluids have been known to be more efficient and environmental friendly extraction fluids (Rodríguez et al. 2014).

Cocoa butter from cocoa bean was extracted using SFE technology by many researchers (McHugh and Krukoniš 1994; Li and Hartland 1996; Saldaña et al. 2002; Tan et al. 2008; Asep et al. 2008, 2013; Jinap et al. 2013; Rodríguez et al. 2014). Less than 5 % of the cocoa butter from the nibs were extracted at 48.3 MPa for a period of 8 h at 40 to 60 °C (McHugh and Krukoniš 1994). The authors concluded that the extraction of cocoa butter by scCO₂ was time consuming and resulted lower yield. In order to facilitate the extraction process, the authors suggested that the addition of ethanol as a cosolvent may enhance the solubility of cocoa butter in scCO₂ (Rossi et al. 1989; Mchugh and Krukoniš 1994; Li and Hartland 1996; Asep et al. 2013).

In contrast to the numerous patents for coffee decaffeination, only Li and Hartland (1992,1996) have conducted studies on the extraction of cocoa butter by scCO₂ using a cosolvent. Nevertheless, they did not optimize SFE process conditions for cocoa butter extraction. In addition, there are limited data reporting cocoa butter extraction optimization with respect to the effects of flow rate and other variables on the composition of TG and FA in butter (Tan et al. 2008).

With respect to cocoa butter extraction using scCO₂, flow rate was studied to allow cocoa butter to reach equilibrium solubility in scCO₂. The extraction of cocoa butter has been pronounced with scCO₂ flow rate. An enhancement in the recovery and solubility of cocoa butter can be achieved using appropriate pressure and temperature. Machmudah et al. (2006) reported that the extraction yield of carotenoids obtained from the optimal processing of Rosehip fruit using scCO₂ was increased by flow rate, pressure and temperature. As expected, the extraction rate constant increased with increasing pressure and flow rate. In order to develop cocoa butter extraction processing using scCO₂ and to determine the efficiency of extraction process, the variables involved in the extraction process (i.e., flow rate, pressure and temperature of extraction) should be studied. The aim of this study was to investigate the effects of scCO₂ extraction process variables, such as flow rate, pressure, and temperature on the yield of cocoa butter. The study also estimates the solubility and selectivity of TG and FA methyl esters.

Materials and Methods

Materials

Cocoa liquor was purchased from K.L. Kepong Sdn. Bhd., Port Klang, Malaysia. Liquid CO₂ with a purity of 99.9 % was obtained from Malaysian Oxygen (MOX), Petaling Jaya, Selangor, Malaysia. Chromatography grade solvent was used in high pressure liquid chromatography (HPLC) and gas chromatography (GC) analysis. Standard of

triglycerides, fatty acid methyl esters, ethanol, isopropanol, acetone, ethanol, ethylene glycol, petroleum ether, methanol and acetonitrile were purchased from Sigma, Merck, and Fisher (Malaysia).

Supercritical fluid extraction (SFE) method for studying the effects of flow rate, pressure and temperature

This experiment used supercritical fluid extraction with pure CO₂ as a solvent (scCO₂). The SFE apparatus consisted of an Intelligent HPLC pump (model PU-1580, Jasco Corporation, Tokyo, Japan) fitted with a cooling jacket to deliver CO₂. The detailed procedure for SFE process was discussed in our previous report (Asep et al. 2013). The liquid carbon dioxide was compressed to the desired pressure of 20 and 35 MPa and then continuously pumped through the extractor at 0.5 (f1), 1.0 (f2), 2.0 (f3) and 4.0 (f4) mL min⁻¹ of flow rates.

The volume of gas that flowed through the system was calculated by the following equation:

$$\text{Weight of CO}_2\text{ gas used (g)} = t \times f \times \delta \quad (1)$$

where,

t is the extraction time (min),

f is the CO₂ gas flow rate (mL min⁻¹), and

δ is the density of CO₂ gas at 20 and 35 MPa and 60 °C (g/mL).

Determination of yield

The determination of the total yield of the extracted cocoa butter was carried out according to the method described in our previous report (Asep et al. 2013).

Determination of triglycerides (TG)

Triglycerides composition was determined by HPLC according to the AOCS method (1993). The detailed procedure for triglycerides analysis was discussed in our previous report (Asep et al. 2013).

Determination of fatty acid methyl esters (FAMES)

The fatty acids (FA) composition of fat mixtures was determined as fatty acid methyl esters (FAMES) using gas chromatography and Restek Rtx-2330 column (30 m; 0.25 mm; 0.2 μm). The detailed procedure for fatty acid analysis was discussed in our previous report (Asep et al. 2013).

Selectivity (α)

The effects of scCO₂ process variables such as flow rate, pressure, and temperature on TG profile and fatty acid (FA)

composition of cocoa butter extracted was determined by estimating the selectivity (α) as function of extraction time. As shown in the following equation, the selectivity was calculated based on the mass balance of TG and FA concentration in the sample and the extract phase:

$$(\alpha) = \frac{[(X_1)_{Ex}] / [(X_1)_S]}{[(X_2)_{Ex}] / [(X_2)_S]} \quad (3)$$

where $(X_1)_{Ex}$ and $(X_2)_{Ex}$ denote as concentration of two different TGs or FAs in the extract phase (supercritical fluid); while $(X_1)_S$ and $(X_2)_S$ are considered as concentration of two different TGs or FAs in the solid phase (soxhlet extraction) (Li and Hartland 1992; AOCS 1993; Rossi 1996). In this research, the selectivity (α) of TG was calculated by comparison of TG relative to 1-palmitoyl-3-stearoyl-2-oleoyl-glycerol (POS) (C52) as the major TG of cocoa butter. Similarly, the selectivity (α) of FA was calculated by comparison of FA relative to stearic acid (C18:0) as the major FA of cocoa butter.

Statistical analysis

Statistical analysis was carried out using the Statistical Analysis System (SAS 1999) Version 8 (TS M1, SAS Institute Inc., Cary, NC, USA) for analysis of variance (ANOVA) and significant differences between means for all treatments (Duncan's multiple range test) at a level of $P < 0.05$.

Results and Discussion

Effects of flow rate, pressure, and temperature on cocoa butter yield

In an attempt to investigate the effect of solvent flow rate on cocoa butter extraction from cocoa liquor, a set of experiments was carried out at 50 and 60 °C temperature, 20 and 35 MPa pressure and the flow rate of supercritical carbon dioxide (scCO₂) was varied at 0.5 (f1), 1.0 (f2), 2.0 (f3) and 4.0 (f4) mL min⁻¹. Figures 1(a), 2(a), 3(a) and 4(a) show the

Fig. 1 Effect of flow rate on cocoa butter extraction from cocoa liquor using scCO₂ at 20 MPa and 50 °C, **a** yield (wt%) versus extraction time (h), **b** yield (wt%) versus weight of CO₂(g). Standard deviations of the reported results are in the range of 0.08 to 1.21, 0.07 to 1.06, 0.09 to 1.14 and 0.09 to 1.11 for f1, f2, f3 and f4 curves, respectively





Fig. 2 Effect of flow rate on cocoa butter extraction from cocoa liquor using scCO₂ at 35 MPa and 50 °C, a yield (wt%) versus extraction time (h), b yield (wt%) versus weight of CO₂(g). Standard deviations of the reported results are in the range of 0.09 to 1.40, 0.07 to 1.04, 0.10 to 1.18 and 0.08 to 1.21 for f1, f2, f3 and f4 curves, respectively




plots for the effect of the scCO₂ flow rate on the yield of cocoa butter extracted as a function of extraction time at 20 MPa, 35 MPa, 50 °C and 60 °C, respectively. It can be observed that cocoa butter yield significantly increased with the CO₂ flow rate. At 20 and 35 MPa, a flow rate of 4.0 mL min⁻¹ (f4) showed the highest yield, followed by 2.0, 1.0 and 0.5 mL min⁻¹. At 20 MPa, 60 °C and 20 h extraction time, a flow rate of 4.0 mL min⁻¹ resulted in the highest yield of 54.80 %, followed by 2.0, 1.0 and 0.5 mL min⁻¹ yielding 42.23, 27.76 and 22.48 % of cocoa butter, respectively. An increase in pressure up to 35 MPa at 60 °C, resulted in significant increase in the yield at 20 h extraction time. A flow rate of 4.0 mL min⁻¹ resulted in nearly 100 % yield, followed by 2.0, 1.0 and 0.5 mL min⁻¹, which produced yields of 89.40, 66.22 and 50.88 %, respectively. Therefore, it can be observed that cocoa butter extraction yield increases with scCO₂ pressure. In addition, it can be observed from the results that cocoa butter yield significantly increased with increase in extraction temperature. The scCO₂ extraction process at

the 60 °C produced a higher cocoa butter yield compared to 50 °C. Jun-ping et al. (2012) reported that the extraction yield of seed oil from *Nigella glandulifera* Freyn using scCO₂ was increased with increasing flow rate, pressure and temperature.

Figures 1(b), 2(b), 3(b) and 4(b) show the effect of the flow rate on cocoa butter yield as a function of the amount of CO₂ used at pressures of 20 and 35 MPa and temperatures of 50 and 60 °C, respectively. Figures 1(b) and 3(b) show that at 20 MPa, the amount of cocoa butter extracted significantly decreased with CO₂ flow rate using the same amount of CO₂. At 20 MPa and 60 °C, when 1000 g CO₂ was used for cocoa butter extraction, the highest amount of the extract obtained was 47.08 % at a flow rate 0.5 mL min⁻¹, followed by 32.32, 26.36 and 18.16 % yield at 1.0, 2.0 and 4.0 mL min⁻¹ flow rate, respectively. Furthermore, Fig. 4(b) shows that the use of 1000 g CO₂ for the extraction of cocoa butter at 35 MPa and 60 °C, resulted in higher yields as compared to those under other

Fig. 3 Effect of flow rate on cocoa butter extraction from cocoa liquor using scCO₂ at 20 MPa and 60 °C, **a** yield (wt%) versus extraction time (h), **b** yield (wt%) versus weight of CO₂(g). Standard deviations of the reported results are in the range of 0.08 to 1.27, 0.06 to 1.09, 0.10 to 1.44 and 0.09 to 1.10 for f1, f2, f3 and f4 curves, respectively



conditions (20 MPa, 50 °C; 20 MPa, 60 °C; 35 MPa, 50 °C), with the highest amount of extracts obtained being 76.04 % at flow rate of 0.5 mL min⁻¹, followed by 67.06, 64.73, 46.88 % at 2.0, 1.0 and 4.0 mL min⁻¹, respectively

On the other hand, from Fig. 4(b), it can be observed that a 70 % yield was obtained when 837.5 g of CO₂ was used at a flow rate of 0.5 mL min⁻¹, followed by 1158.4, 1075.1, and 1595.2 g at 1.0, 2.0, and 4.0 mL min⁻¹, respectively. A flow rate of 0.5 mL min⁻¹ consumed less CO₂ than the other flow rates. CO₂ consumption at a flow rate of 2.0 mL min⁻¹ was very close to that at a flow rate of 1.0 mL min⁻¹. Interestingly, the graph of the optimum flow rate of 2.0 mL min⁻¹ at 35 Mpa was observed to overlap (crossover) with that of 1.0 mL min⁻¹ flow rate; indicating that the equilibrium solubility limit had been achieved, and the interaction between solute and solvent may have been optimum.


From Fig. 4(b), we can see that the flow rate of 2.0 mL min⁻¹ shows a better yield than 1.0 and 4.0 mL min⁻¹. Although 0.5 mL min⁻¹ resulted in the highest yield with a consumption of 1000 g CO₂, it required 38.5 h;

thus, the extraction time used was longer as compared to 1.0 and 2.0 mL min⁻¹ flow rates, which required extraction time of 19.5 and 9.5 h, respectively. Although 4.0 mL min⁻¹ showed the highest yield, the yield can be decreased by the residence time of CO₂ in the extraction vessel, and therefore, the time of contact between the solvent and sample was shorter; however, 4.0 mL min⁻¹ require more CO₂ (1595.2 g CO₂ at 70 % yield) for extraction. Therefore, 2.0 mL min⁻¹ was considered the optimum CO₂ flow rate that can provide the greatest effect on the cocoa butter extraction performance.

Effects of flow rate, pressure, and temperature on triacylglycerol

Triacylglycerols (TGs) are the major component present in cocoa butter, making up 92 to 96 % of the lipid composition (Jahurul et al. 2013). Tables 1 and 2 present the results of the TG analysis of cocoa butter obtained at 0.5 to 4.0 mL min⁻¹ flow rate, 20 and 35 MPa pressure and 50 and 60 °C

Fig. 4 Effect of flow rate on cocoa butter extraction from cocoa liquor using scCO₂ at 35 MPa and 60 °C, a yield (wt%) versus extraction time (h), b yield (wt%) versus weight of CO₂(g). Standard deviations of the reported results are in the range of 0.09 to 1.43, 0.09 to 1.27, 0.11 to 1.55 and 0.07 to 1.19 for f1, f2, f3 and f4 curves, respectively



temperature using scCO₂. The results show that cocoa butter extracted from various treatments had three main TGs; POS, 1,3-dipalmitoyl-2-oleoyl-glycerol (POP) and 1,3-distearoyl-2-oleoyl-glycerol (SOS). The percentage of each TG composition was significantly different ($P < 0.05$) among various flow rates and pressures, except for POS. POS was identified as the major TG component in 42.26 to 45.12 % cocoa butter yields, followed by SOS and POP with 27.05 to 30.67 % and 21.05 to 23.64 %, respectively. These findings were in good agreement with previously reported data on the typical TG composition of cocoa butter (Chaiseri and Dimick 1989; Lipp et al. 2001; Spangenberg and Dionisi 2001; Ali et al. 2001; Tan et al. 2008; Asep et al. 2008, 2013; Jinap et al. 2013). Cocoa butter contains high amounts of symmetrical monounsaturated TG, in particular 1,3-dipalmitoyl-2-oleoylglycerol (POP), 1-palmitoyl-2-oleoyl-3-stearoyl-glycerol (POS) and 1,3-distearoyl-2-oleoyl-glycerol (SOS). These TGs are of particular interest, since they are mainly responsible for the desired properties of chocolate, i.e. crystallization and melting behavior.

As can be seen in Tables 1 and 2, at the same pressure and temperature, POP and POS generally decrease with flow rate and extraction time; in contrast, SOS increase with flow rate and extraction time. When the pressure and temperature are higher, POP and POS decrease with an increase in flow rate and extraction time; however, SOS increase with flow rate and extraction time. In general, POP, POS and SOS showed higher solubility at a higher pressure of 35 MPa and a temperature of 60 °C compared to 20 MPa and 50 °C. This result may be explained by the fact that the increase in pressure and temperature cause an increase the TG solubility. These results show the progressive extraction of cocoa butter; the more soluble POP triglyceride (MW slightly smaller) is easily extracted in the early stages, followed by POS, and the less soluble SOS (MW slightly larger) is removed in the late stages. This behavior is supported by Rossi et al. (1989) who also investigated the composition of cocoa butter by scCO₂ of various cocoa products and, observed minor differences in the

Table 1 Triacylglycerol composition (area %)^α of cocoa butter extracted by supercritical carbon dioxide (scCO₂) for effect of flow rate (F) with different pressures at 50 °C

Sample ^β	5 h Extraction Time					10 h Extraction Time					15 h Extraction Time				
	POP	POS	SOS	Others ^γ	Others ^γ	POP	POS	SOS	Others ^γ	Others ^γ	POP	POS	SOS	Others ^γ	
	F1 (0.5 mL min ⁻¹ ; 20 MPa)	22.75 ± 0.33	44.31 ± 0.54	27.99 ± 0.70	4.95 ± 0.06	22.67 ± 0.39	44.10 ± 0.65	28.36 ± 0.33	4.87 ± 0.06	22.89 ± 0.31	42.64 ± 0.56	29.67 ± 0.54	4.80 ± 0.04		
F2 (1.0 mL min ⁻¹ ; 20 MPa)	22.34 ± 0.41	44.21 ± 0.53	28.61 ± 1.15	4.83 ± 0.13	22.13 ± 0.43	44.01 ± 0.62	29.10 ± 0.51	4.76 ± 0.14	22.87 ± 0.82	42.35 ± 0.53	30.10 ± 0.78	4.69 ± 0.09			
F3 (2.0 mL min ⁻¹ ; 20 MPa)	22.47 ± 0.93	44.13 ± 0.97	28.53 ± 0.69	4.86 ± 0.43	22.07 ± 0.92	43.67 ± 1.08	29.48 ± 0.31	4.78 ± 0.44	22.43 ± 1.68	42.26 ± 0.87	30.60 ± 0.43	4.71 ± 0.25			
F4 (4.0 mL min ⁻¹ ; 20 MPa)	22.01 ± 1.35	44.08 ± 1.20	28.61 ± 0.52	5.30 ± 0.23	21.74 ± 1.49	43.36 ± 1.49	29.96 ± 0.38	4.94 ± 0.27	22.19 ± 2.78	42.28 ± 1.34	30.67 ± 0.61	4.87 ± 0.17			
F5 (0.5 mL min ⁻¹ ; 35 MPa)	22.86 ± 1.34	44.64 ± 1.33	27.66 ± 0.68	4.84 ± 0.68	22.85 ± 1.79	44.27 ± 1.80	28.11 ± 0.61	4.77 ± 0.95	22.78 ± 2.59	44.03 ± 1.95	28.50 ± 1.17	4.69 ± 0.40			
F6 (1.0 mL min ⁻¹ ; 35 MPa)	23.01 ± 1.49	44.20 ± 1.40	27.85 ± 1.47	4.94 ± 0.53	22.71 ± 1.29	44.06 ± 1.23	28.37 ± 0.85	4.86 ± 0.48	22.64 ± 1.22	43.99 ± 0.87	28.58 ± 1.07	4.79 ± 0.13			
F7 (2.0 mL min ⁻¹ ; 35 MPa)	23.20 ± 1.32	43.65 ± 0.92	28.25 ± 1.17	4.90 ± 0.72	22.45 ± 1.55	43.96 ± 1.09	28.76 ± 0.92	4.83 ± 0.87	22.46 ± 1.83	43.96 ± 1.04	28.82 ± 1.56	4.76 ± 0.33			
F8 (4.0 mL min ⁻¹ ; 35 MPa)	23.64 ± 0.20	42.56 ± 0.40	28.73 ± 0.65	5.07 ± 0.05	22.04 ± 0.19	43.77 ± 0.39	29.19 ± 0.26	5.00 ± 0.05	22.43 ± 0.18	43.51 ± 0.40	29.14 ± 0.27	4.93 ± 0.05			

^α Means value ± standard deviation of three replications

^β Cocoa butter extracted obtained by scCO₂ at 50 °C with different flow rate and pressure

^γ PLIO, PLP, POO, SOO, and SOA where P = palmitic, O = oleic, S = stearic, Li = linoleic, A = arachidic

selectivity of the process for TG of cocoa butter, namely, POP (C50), POS (C52) and SOS (C54). At 50 °C and 40 MPa, the average selectivity of POP relative to POS was 1.05, whereas that of SOS relative to POS was 0.94. Consequently, Rossi et al. (1989) hypothesized that shorter TGs are more easily extracted than longer ones.

Ragunath et al. (1992, 1993), Hassan et al. (2000, 2001), Markom et al. (2001), Norulaini et al. (2004) and Zaidul et al. (2006) fractionated triglycerides of fatty oil components based on their carbon number using scCO₂ at a temperature range of 40 to 80 °C and pressures up to 50 MPa. Researchers reported a low solubility of TGs rich crude palm oil in scCO₂ at relatively low temperature and low pressure while at any temperature and pressure the short chain fatty acids are eluted fast followed by medium and long chain.

Arul et al. (1994) reported that the solubilities of short-, medium- and long chain TGs increased with pressure, temperature and flow rate of CO₂. Another important factor that must be considered is the pressure dependence of the dissolving power of supercritical fluids (Medir and Giralt 1982; Bling and Franck 1983). At constant temperature the volatility or fugacity of the TGs phase increases with pressure. Perry et al. (1949) reported that the vapor pressure of TG decreases with their carbon number. Therefore, the short chain TGs are, in general, more volatile than the long chain molecules. The concentration of long-chain TG is higher than the short chain in milk fat at scCO₂ reported by Arul et al. (1994).

Effects of flow rate, pressure, and temperature on fatty acid methyl ester

The results of fatty acid methyl esters obtained from gas chromatography analysis are shown in Tables 3 and 4. Palmitic acid (C16:0), stearic acid (C18:0) and oleic acid (C18:1) where the three main fatty acids in the cocoa butter extracted using scCO₂, with C18:0 of 32.53 to 38.92 % being the major component, followed by C18:1 with 30.62 to 33.06 % and C16:0 with 25.94 to 31.49 %. It can be seen from Tables 3 and 4 that the FA compositions of cocoa butter were significantly affected (*P* < 0.05) by flow rate with different pressures and temperatures. These findings are in good agreement with the typical FA compositions of cocoa butter obtained by Chaiseri and Dimick (1989), Lipp et al. (2001), Spangenberg and Dionisi (2001), Ali et al. (2001), Tan et al. (2008), Asep et al. (2008, 2013) and Jinap et al. (2013).

In addition, at the same pressure, C16:0 and C18:2 generally decreased with an increase in flow rate and extraction time, whereas C18:0 and C18:1 increased with the increase in flow rate and decreased with an increase in extraction time.

The results also show that when the pressure increased, in general C16:0 decreased with an increase in flow rate and

Table 2 Triacylglycerol composition (area %)^α of cocoa butter extracted by supercritical carbon dioxide (scCO₂) for effect of flow rate (f) with different pressures at 60 °C

Sample ^β	5 h Extraction Time				10 h Extraction Time				15 h Extraction Time			
	POP	POS	SOS	Others ^γ	POP	POS	SOS	Others ^γ	POP	POS	SOS	Others ^γ
f1 (0.5 mL min ⁻¹ ; 20 MPa)	22.83 ±0.25	44.70 ±0.56	27.54 ±0.65	4.93 ±0.07	23.41 ±0.37	44.12 ±0.45	27.07 ±0.35	4.88 ±0.07	23.12 ±0.21	43.96 ±0.45	28.09 ±0.38	4.83 0.05
f2 (1.0 mL min ⁻¹ ; 20 MPa)	22.80 ±0.31	44.66 ±0.54	27.72 ±1.22	4.82 ±0.19	23.22 ±0.43	43.97 ±0.41	28.04 ±0.62	4.77 ±0.19	23.10 ±0.55	43.66 ±0.41	28.52 ±0.65	4.72 0.13
f3 (2.0 mL min ⁻¹ ; 20 MPa)	22.70 ±0.77	44.58 ±1.00	27.87 ±0.73	4.85 ±0.60	22.89 ±1.02	43.82 ±0.72	28.48 ±0.37	4.80 ±0.57	22.43 ±1.22	43.12 ±0.67	29.69 ±0.37	4.75 0.39
f4 (4.0 mL min ⁻¹ ; 20 MPa)	22.01 ±1.11	44.33 ±1.23	28.36 ±0.55	5.30 ±0.33	21.64 ±1.65	43.58 ±0.99	29.82 ±0.59	4.97 ±0.34	21.97 ±2.02	42.70 ±1.03	30.41 ±0.65	4.92 0.26
f5 (0.5 mL min ⁻¹ ; 35 MPa)	22.98 ±1.10	45.12 ±1.57	27.05 ±0.72	4.85 ±0.96	23.47 ±1.98	44.41 ±1.53	27.32 ±0.94	4.80 ±1.22	23.24 ±1.89	44.12 ±1.93	27.89 ±1.26	4.75 0.63
f6 (1.0 mL min ⁻¹ ; 35 MPa)	22.91 ±1.36	44.75 ±1.65	27.39 ±1.55	4.95 ±0.75	22.75 ±1.58	44.44 ±1.04	27.91 ±1.31	4.90 ±0.62	22.52 ±0.98	43.99 ±0.86	28.63 ±1.14	4.85 0.21
f7 (2.0 mL min ⁻¹ ; 35 MPa)	22.85 ±1.21	44.70 ±1.09	27.52 ±1.24	4.92 ±1.01	22.69 ±1.90	44.39 ±0.93	28.05 ±1.42	4.87 ±1.13	22.47 ±1.47	43.94 ±1.03	28.77 ±1.67	4.82 0.51
f8 (4.0 mL min ⁻¹ ; 35 MPa)	21.05 ±0.20	44.24 ±0.40	29.61 ±0.65	5.10 ±0.05	21.61 ±0.19	43.93 ±0.39	29.41 ±0.26	5.05 ±0.05	21.88 ±0.18	43.49 ±0.40	29.63 ±0.27	5.00 0.05

^α Means value ± standard deviation of three replications

^β Cocoa butter extracted obtained by scCO₂ at 60 °C with different flow rate and pressure

^γ PLiO, PLiP, POO, SOO, and SOA where P = palmitic, O = oleic, S = stearic, Li = linoleic, A = arachidic

extraction time; however, C18:1 and C18:2 in general increased with an increase in flow rate and a decrease in extraction time; and the C18:0 percentage increased with an increase in flow rate and extraction time. It is interesting to note that the solubility of FAs increased with an increase in pressure and temperature. At a pressure of 35 MPa, the solubility of FA with larger MW is higher because higher pressure provides greater solvating power and enhances the interaction between CO₂ solvents with cocoa butter.

On the other hand, based on the solubility of FA in scCO₂, it was found that at lower pressures (20 MPa), the short chain fatty acid constituents (C16) were extracted with higher selectivity than the long chain fatty acid constituents (C18:0 and C18:1). At higher pressures (35 MPa), the C16:0 started to reduce, and the C18:0 and C18:1 constituents increased. This indicates that more of the shorter chain fatty acid (C16:0) was extracted than the long chain fatty acid constituents (C18:0 and C18:1) at lower pressures (20 MPa), whereas more of the longer chain fatty acids (C18:0 and C18:1) were extracted than the C16:0 at higher pressures. Stahl et al. (1988) reported that the solubility of fatty acids in scCO₂ extraction depends on the length of the hydrocarbon chain and the presence of functional groups. Moreover, it also depends on the effect of the extraction parameters, pressure and temperature (Stahl et al. 1988). This behavior is similar to the findings of previous works investigating the extraction and fractionation of

fatty acids from palm kernel oil (PKO) in scCO₂, as reported by Norulaini et al. (2004) and Zaidul et al. (2006, 2007a, b).

Conclusion

The effects of different flow rate, pressures and temperatures on cocoa butter extraction using scCO₂ were investigated. The results indicated that the extraction yield is significantly ($P < 0.05$) influenced by the variables and the yields was increased with flow rate, pressure, and temperature where the optimum values were 2 mLmin⁻¹, 35 MPa and 60 °C, respectively. The solubility of FAs and TGs were also increased with pressure and temperature. TG and FA compositions were found to be similar to those of cocoa butter obtained by conventional methods. In terms of the selectivity, the lower molecular weight TGs and FAs showed higher selectivity compared to higher molecular weight TGs and FAs. This indicates that POP was more easily extracted at the beginning of the extraction process, while the amount of POS, followed by that of SOS, was found to be higher at the end of the extraction process. In contrast, C16:0 and C18:0 were separated more easily than C18:1, and C18:2 during extraction. Thus, they were fractionated during the first stage of the scCO₂ extraction process.

Table 3 Fatty acids composition (area %)^α changes of cocoa butter extracted by supercritical carbon dioxide (scCO₂) for effect of flow rate (f) with different pressures at 50 °C

Sample ^β	5 h Extraction Time					10 h Extraction Time					15 h Extraction Time				
	C16:0	C18:0	C18:1	C18:2	Others ^γ	C16:0	C18:0	C18:1	C18:2	Others ^γ	C16:0	C18:0	C18:1	C18:2	Others ^γ
f1 (0.5 mL min ⁻¹ ; 20 MPa)	29.08 ±0.35	36.51 ±0.64	30.72 ±0.64	2.40 ±0.11	1.29 ±0.07	28.64 ±0.41	36.57 ±0.61	31.15 ±0.79	2.37 ±0.23	1.27 ±0.08	26.49 ±0.27	38.60 ±0.86	31.54 ±0.47	2.19 ±0.04	1.18 ±0.10
f2 (1.0 mL min ⁻¹ ; 20 MPa)	28.55 ±0.43	36.66 ±0.62	31.38 ±1.20	2.29 ±0.29	1.13 ±0.26	28.33 ±0.48	36.68 ±0.56	31.63 ±1.40	2.25 ±0.58	1.11 ±0.28	26.01 ±0.71	38.87 ±0.78	32.01 ±0.81	2.09 ±0.09	1.03 ±0.23
f3 (2.0 mL min ⁻¹ ; 20 MPa)	28.32 ±0.90	36.84 ±1.14	31.55 ±0.72	2.14 ±0.95	1.15 ±0.98	28.14 ±0.96	36.75 ±0.98	31.87 ±0.80	2.11 ±1.79	1.13 ±0.80	26.12 ±1.17	38.77 ±0.42	32.12 ±1.54	1.95 ±0.24	1.05 ±0.36
f4 (4.0 mL min ⁻¹ ; 20 MPa)	28.11 ±1.31	36.98 ±1.40	31.86 ±0.54	1.97 ±0.51	1.08 ±0.92	28.05 ±1.16	36.84 ±1.15	32.11 ±0.57	1.94 ±0.92	1.06 ±0.72	25.94 ±1.39	38.92 ±0.34	32.36 ±0.88	1.80 ±0.24	0.98 ±0.47
f5 (0.5 mL min ⁻¹ ; 35 MPa)	31.33 ±1.17	33.11 ±1.80	31.05 ±0.71	2.83 ±1.23	1.68 ±1.12	31.18 ±0.98	32.74 ±1.40	31.60 ±0.71	2.81 ±2.10	1.67 ±0.83	30.09 ±1.32	33.44 ±0.80	32.15 ±1.09	2.71 ±0.24	1.61 ±0.54
f6 (1.0 mL min ⁻¹ ; 35 MPa)	30.14 ±1.19	34.37 ±1.89	31.43 ±1.53	2.57 ±0.96	1.49 ±0.87	29.99 ±0.95	33.99 ±1.39	31.98 ±0.86	2.56 ±1.56	1.48 ±0.61	28.94 ±0.85	34.63 ±0.63	32.53 ±0.52	2.47 ±0.12	1.43 ±0.23
f7 (2.0 mL min ⁻¹ ; 35 MPa)	29.82 ±1.06	34.67 ±1.25	31.51 ±1.22	2.49 ±1.30	1.50 ±0.92	29.67 ±0.80	34.30 ±0.87	32.06 ±0.65	2.48 ±2.00	1.50 ±0.61	28.63 ±0.67	34.93 ±0.64	32.61 ±0.91	2.39 ±0.16	1.44 ±0.24
f8 (4.0 mL min ⁻¹ ; 35 MPa)	29.43 ±0.94	34.84 ±0.82	31.74 ±0.98	2.49 ±1.75	1.50 ±0.96	29.28 ±0.68	34.46 ±0.55	32.28 ±0.50	2.48 ±2.56	1.50 ±0.61	28.26 ±0.53	35.08 ±0.66	32.83 ±1.12	2.39 ±0.21	1.44 ±0.26

^α Means value ± standard deviation of three replications

^β Cocoa butter extracted obtained by scCO₂ at 50 °C with different flow rate and pressure

^γ C12:0, C14:0 and C18:3

Table 4 Fatty acids composition (area %)^α changes of cocoa butter extracted by supercritical carbon dioxide (scCO₂) for effect of flow rate (f) with different pressures at 60 °C

Sample ^β	5 h Extraction Time				10 h Extraction Time				15 h Extraction Time				
	C16:0	C18:0	C18:1	Others ^γ	C16:0	C18:0	C18:1	Others ^γ	C16:0	C18:0	C18:1	C18:2	Others ^γ
f1 (0.5 mL min ⁻¹ ; 20 MPa)	29.67 ±0.24	35.94 ±0.63	30.62 ±0.57	1.32 ±0.05	29.23 ±0.36	36.19 ±0.51	30.87 ±0.74	2.41 ±0.09	27.03 ±0.32	38.41 ±0.99	31.12 ±0.24	2.23 ±0.05	1.20 ±0.02
f2 (1.0 mL min ⁻¹ ; 20 MPa)	29.13 ±0.30	36.07 ±0.61	31.32 ±1.07	1.15 ±0.16	28.69 ±0.42	36.31 ±0.47	31.57 ±1.31	2.30 ±0.23	26.54 ±0.84	38.47 ±0.90	31.82 ±0.41	2.13 ±0.12	1.05 ±0.05
f3 (2.0 mL min ⁻¹ ; 20 MPa)	28.96 ±0.74	36.10 ±1.12	31.62 ±0.64	1.16 ±0.63	28.52 ±1.00	36.34 ±0.82	31.87 ±0.75	2.13 ±0.70	26.38 ±1.38	38.47 ±0.48	32.12 ±1.20	1.97 ±0.43	1.06 ±0.08
f4 (4.0 mL min ⁻¹ ; 20 MPa)	28.76 ±1.07	36.30 ±1.38	31.86 ±0.48	1.09 ±0.59	28.33 ±1.38	36.53 ±0.96	32.11 ±0.53	1.96 ±0.36	26.20 ±1.65	38.63 ±0.39	32.36 ±0.69	1.81 ±0.43	0.99 ±0.11
f5 (0.5 mL min ⁻¹ ; 35 MPa)	31.49 ±1.06	32.90 ±1.77	31.08 ±0.63	1.69 ±1.69	31.33 ±1.29	32.53 ±1.17	31.63 ±0.67	2.83 ±1.00	30.24 ±1.56	33.23 ±0.92	32.18 ±1.50	2.73 ±1.02	1.62 ±0.12
f6 (1.0 mL min ⁻¹ ; 35 MPa)	30.14 ±1.30	34.30 ±1.86	31.50 ±1.36	1.49 ±1.32	29.99 ±1.51	33.93 ±1.16	32.05 ±1.37	2.56 ±0.74	28.94 ±1.01	34.57 ±1.23	32.60 ±0.72	2.47 ±0.49	1.43 ±0.05
f7 (2.0 mL min ⁻¹ ; 35 MPa)	29.52 ±1.16	34.85 ±1.23	31.67 ±1.09	1.49 ±1.38	29.38 ±1.28	34.47 ±0.73	32.22 ±1.04	2.45 ±0.95	28.35 ±0.79	35.08 ±1.26	32.77 ±1.25	2.37 ±0.66	1.43 ±0.05
f8 (4.0 mL min ⁻¹ ; 35 MPa)	29.14 ±0.54	34.99 ±0.45	31.96 ±0.36	1.47 ±0.08	28.99 ±0.26	34.60 ±0.36	32.51 ±0.25	2.43 ±0.02	27.98 ±0.26	35.20 ±0.34	33.06 ±0.26	2.34 ±0.02	1.41 ±0.01

^α Means value ± standard deviation of three replications^β Cocoa butter extracted by scCO₂ at 60 °C with different flow rate and pressure^γ C12:0, C14:0 and C18:3

Acknowledgments The authors would like to thank to the Ministry of Science, Technology and Environment of Malaysia for supporting financially to this research under Intensification of Research Priority Area (IRPA) Project No. 01-02-04-0512.

References

- Ali A, Selamat J, Che Man YB, Suria AM (2001) Characterization and fat migration of palm kernel stearin as affected by addition of desiccated coconut used as base filling Centre in dark chocolate. *Int J Food Sci Nutr* 52:251–261
- American Oil Chemist Society (AOCS 1993) Official and tentative methods of American oil chemist society, fourth ed., American Oil Chemist Society, Champaign, USA, pp1-5.
- Arul J, Tardif R, Boudreau A, Douglas ST, McGinnis LRW (1994) Solubility of milk fat triglycerides in supercritical carbon dioxide. *Food Res Int* 21:459–467
- Asep EK, Jinap S, Tan TJ, Russly AR, Harcharan S, Nazimah SAH (2008) The effects of particle size, fermentation and roasting of cocoa nibs on supercritical fluid extraction of cocoa butter. *J Food Eng* 85:450–458
- Asep EK, Jinap S, Jahurul MHA, Zaidul ISM, Singh H (2013) Effects of polar cosolvents on cocoa butter extraction using supercritical carbon dioxide. *Innov Food Sci Emerg Technol* 20:152–160
- Bling RGL, Franck EU (1983) Calculation of phase equilibria for separation operations of *Ber*. *Bunsenges J Phys Chem* 87:882–890
- Campardelli R, Adami R, Porta GD, Reverchon E (2012) Nanoparticle precipitation by supercritical assisted injection in a liquid antisolvent. *Chem Eng J* 192:246–251
- Chaiseri S, Dimick PS (1989) Lipid and hardness characteristics of cocoa butters from different geographic regions. *J Am Oil Chem Soc* 66:1771–1776
- Hassan MN, Rahman NNA, Anuar BO, Ibrahim MH, Mohd Omar AK (2000) Simple fractionation through the supercritical carbon dioxide extraction of palm kernel oil. *Sep Purif Technol* 19:113–120
- Hassan MN, Rahman NNA, Anuar BO, Ibrahim MH, Mohd Omar AK (2001) The effect of dehulling on supercritical extraction of palm kernel oil. *J Chem Eng Japan* 34:407–410
- Jahurul MHA, Zaidul ISM, Sahena F, Yazid AMM, Norulaini NAN, Omar AKM (2012) Application of supercritical fluid extraction (SFE) of palm oil and oil from natural sources. *Molecules* 17:1764–1794
- Jahurul MHA, Norulaini NAN, Zaidul ISM, Jinap S, Sahena F, Mohd Omar AK (2013) Cocoa butter fats and possibilities of substitution in food products concerning cocoa varieties, alternative sources, extraction methods, composition, and characteristics. *J Food Eng* 117(4):467–476
- Jahurul MHA, Zaidul ISM, Nik Norulaini NA, Sahena F, Kamaruzzaman BY, Kashif Ghafoor, Omar AKM. (2014). Cocoa butter replacers from blends of mango seed fat extracted by supercritical carbon dioxide and palm stearin. *Food Res Int* 65: 401–406.
- Jahurul MHA, Zaidul ISM, Norulaini NAN, Sahena F, Rahman MM, Mohd Omar AK (2015) Optimization of supercritical carbon dioxide extraction parameters of cocoa butter analogy fat from mango seed kernel oil using response surface methodology. *J Food Sci Technol* 52(1):319–326
- Jinap S, Asep EK, Jahurul MHA, Russly AR, Sahena F, Zaidul ISM (2013) Effects of moisture and pH on supercritical fluid extraction of cocoa butter. *Food Bioprocess Technol* 6:2455–2465
- Knez Ž, Škerget M, Knez Hrnčič M, Čuček D (2014) Particle formation using sub- and supercritical fluids. In: Anikeev V, Fan M (eds) *Supercritical fluid technology for energy and environmental applications*. Boston, pp. 31–67
- Li S, Hartland S (1992) Influence of co-solvents on solubility and selectivity in extraction of xanthines and cocoa butter from cocoa beans with supercritical CO₂. *J Supercrit Fluid* 5:7–12
- Li S, Hartland S (1996) A new industrial process for extracting cocoa butter and xanthines with supercritical carbon dioxide. *J Am Oil Chem Soc* 73(4):423–429
- Lipp M, Simoneau C, Ulberth F, Anklam E, Crews C, Brereton P, de Greyt W, Schwack W, Wiedmaier C (2001) Composition of genuine cocoa butter and cocoa butter equivalents. *J Food Compos Anal* 14:399–408
- Machmudah S, Shotipruk A, Goto M, Sasaki M, Hirose T (2006) Extraction of astaxanthin from *Haematococcus Pluvialis* using supercritical CO₂ and ethanol as entrainer. *Ind Eng Chem Res* 45:3652–3657
- Markom M, Singh H, Hasan M (2001) Supercritical CO₂ fractionation of crude palm oil. *J Supercrit Fluid* 20:45–53
- Mchugh MA, Krukonis VJ (1994) *Supercritical fluid extraction: principles and practice*, second ed., Boston, Butterworth.
- Medir M, Giralt F (1982) Correlation of activity coefficients of hydrocarbons in water at infinite dilution with molecular parameters. *AIChE J* 28:341–343
- Norulaini NNA, Zaidul ISM, Anuar O, Mohd Omar AK (2004) Supercritical enhancement for separation of lauric acid and oleic acid in palm kernel oil (PKO). *Sep Purif Technol* 35(1):55–60
- Perry ES, Weber WH, Daubert BF (1949) Vapor pressure of phlegmatic liquids: I. simple and mixed triglycerides. *J Am Oil Chem Soc* 71:3720–3726
- Ragunath B, Inomata H, Adschiri T, Arai K (1992) Phase equilibrium study for the separation and fractionation of fatty oil components using supercritical carbon dioxide. *Fluid Phase Equilib* 81:307–320
- Ragunath B, Yamane S, Inomata H, Adschiri T, Arai K (1993) Phase equilibria of supercritical CO₂ fatty oil component binary systems. *Fluid Phase Equilib* 83:183–192
- Reverchon E, Adami R (2006) Nanomaterials and supercritical fluids. *J Supercrit Fluid* 37:1–22
- Rodríguez Ó, Ortuño C, Simal S, Benedito J, Femenia A, Rosselló C (2014) Acoustically assisted supercritical CO₂ extraction of cocoa butter: effects on kinetics and quality. *J Supercrit Fluid* 94:30–37
- Rossi M, Arnoldi C, Salvioni G, Schiraldi A (1989) Characterization of cocoa extracts obtained with supercritical carbon dioxide. *Italian J Food Sci* 3:41–50
- Rossi M (1996) Supercritical fluid extraction of cocoa and cocoa products. In: King JW, List GR (eds) *Supercritical fluid technology in oil and lipid chemistry*. AOCS Press, Champaign, Illinois, pp. 220–229
- Saldaña MDA, Mohamed RS, Mazzafera P (2002) Extraction of cocoa butter from Brazilian cocoa beans using supercritical CO₂ and ethanol. *Fluid Phase Equilib* 194–197:885–894
- Spangenberg JE, Dionisi F (2001) Characterization of cocoa butter and cocoa butter equivalents by bulk and molecular carbon isotope analyses: implications for vegetable fat quantification in chocolate. *J Agric Food Chem* 49:4271–4277
- Statistical Analysis System (SAS 1999). USA: SAS institute Inc., Cary, NC.
- Stahl E, Quirin KW, Gerald D (1988) *Dense gases for extraction and refining*. Springer-Verlag, New York, p. 176
- Tan TJ, Jinap S, Asep EK, Nazimah SHA (2008) Extraction of cocoa butter by supercritical carbon dioxide: optimization of operating conditions and effect of particle size. *J Food Lipid* 15:263–276
- Zaidul ISM, Norulaini NAN, Mohd Omar AK (2006) Separation and fractionation of triglycerides in terms of fatty acid constituents in palm kernel oil using supercritical CO₂. *J Sci Food Agric* 86:1138–1145
- Zaidul ISM, Norulaini NNA, Mohd Omar AK, Smith Jr RL (2007a) Blending of supercritical carbon dioxide (SC-CO₂) extracted palm kernel oil fractions and palm oil to obtain cocoa butter replacers. *J Food Eng* 78:1397–1409
- Zaidul ISM, Norulaini NNA, Mohd Omar AK, Smith Jr RL (2007b) Supercritical carbon dioxide (SC-CO₂) extraction of palm kernel oil from palm kernel. *J Food Eng* 79:1007–1014