

RP 501

CPCRI

Reprinted from Potassium in Soils and Crops, proceedings of a Symposium held at Vigyan Bhavan, New Delhi, Nov. 16-17, 1978. Edited by G. S. Sekhon, published by Potash Research Institute of India, New Delhi.

**K NUTRITION
OF PLANTATION CROPS—
A REVIEW OF WORK DONE
IN INDIA**

E. V. Nelliatt

*Central Plantation Crops Research Institute,
Kasaragod 670124. Kerala*

**POTASH RESEARCH INSTITUTE OF INDIA
NEW DELHI**

K NUTRITION OF PLANTATION CROPS— A REVIEW OF WORK DONE IN INDIA

E.V. NELLIAT

*Central Plantation Crops Research Institute,
Kasaragod 670124, Kerala:*

ABSTRACT

Potash is the nutrient element most consumed by plantation crops. Although response to K differs widely among crops, it generally improves the yield and quality of the products.

Potash deficiency symptoms in various plantation crops are reviewed. In coconut, potassium deficiency leads to chlorosis, leaf scorch and development of poor crown with short fronds. In arecanut, the deficiency symptoms consist of dead tissues around the leaf margin and between the veins of leaves. In cardamom, potassium deficiency results in the browning of leaves and reducing sucker production. In rubber, the characteristic symptom is the development of marginal and tip chlorosis in leaves. In tea, K deficiency leads to defoliation and debility. In coffee, a striking leaf tip and leaf edge blackening develops.

The requirements of coconut, tea, coffee and rubber for potassium range from 220 to 300 kg K_2O per hectare. In coconut adequate K nutrition reduces the pre-bearing age and increase the fruit set, husk weight and nut size. Copra out-turn per palm increases by over 20%. Potash fertilization to cashew and arecanut enhances the fruit set and nut weight. In coffee and tea, K nutrition helps in the development of a strong frame. Little information is available on plantation crops other than coconut, rubber, tea and coffee.

INTRODUCTION

Plantation crops are a group of commercial crops of perennial nature.

They are cultivated extensively in the tropics and the sub-tropics and demand employment of labour throughout the year. The produce of these crops had entered the international trade long before the food grains. Although area under these crops is a mere fraction of the world's cultivated area, their impact on the economy of the concerned region or the country is immense. In India, plantation crops occupy less than 2% of the cultivated area but account for 75% of our export earnings from agricultural commodities and 25% of our earnings from all exports¹. The statistics on area, production, export and import of the major plantation crops are given in Table 1.

A plantation crop occupies the land for decades. Balanced fertilisation commensurate with the crop need and the soil's capacity to supply nutrients, is essential for maintaining a high level of production. It should fully meet the twin requirements viz., replenish the amount of nutrients removed in crop harvests from time to time, and supply enough for the normal growth and physiological functions of the plant. An imbalance in the nutritional status of the crop may be disastrous.

The plantation crops, in general, are heavy potassium feeders and potassium nutrition is a major limiting factor in their productivity. Accordingly, fertiliser recommendations for many of these crops have N : K ratio of 1 : 2. Sufficient information on the economic optima of their nutritional requirements is, however, not available for different agro-climatic regions. The available literature and experimental data on the response of major plantation crops to potassium are reviewed and assessed in this paper.

COCONUT (COCOS NUCIFERA)

Coconut is grown on soil types ranging from littoral sand to heavy clay and pH ranging from 3.5 to 8.9². Most of these soils are low in available nutrients. Coconut is known to be a potash loving plant on account of the abundance of potassium in all parts of the palm, especially the growing portion³. The critical levels of N and K, on dry matter basis, in the standard coconut leaf are 1.8—2.0% and 0.8—1.0% respectively. The nutrient content of the leaves of young coconut palms under NPK fertiliser experiment at CPCRI Kasaragod⁴ are presented in Table 2.

The potassium deficiency symptoms in the initial stages of crop growth are drying up of the leaf tips and development of scattered olive green spots, starting from leaf tips. The spots increase in size and later turn

TABLE 1: Plantation Crops in India—Area, production, exports and imports

Crop	Area ('000ha)	Production ('000 tonnes)	Average yield (kg/ha)	Export (Million Rs.)	Import	Leading states in India and their share (%) of area
Tea	361.4	487.4	1360	1530.9	Nil	Assam 51.8 West Bengal 24.4 Kerala 10.1
Coffee	156.1	92.5	590	584.0	Nil	Karnataka 59.2 Kerala 23.6 Tamil Nadu 16.2
Rubber	225.0	138.0	760	Nil	Nil	Kerala 86.2 Tamil Nadu 5.0 Karnataka 2.6
Coconut	1115.5	5961.3 (million nuts)	5344 (Nuts/ha)	177.1	2.7	Kerala 62.4 Tamil Nadu 16.6 Karnataka 12.7
Arecanut	187.4	169.4	904	5.3	Nil	Karnataka 42.44 Kerala 33.8 Assam 18.9
Cashewnut	418.1	184.7	442	1196.2	366.0	Kerala 63.8 Karnataka 18.1 Andhra Pradesh 6.9
Cacao	3.6	—	—	4.6	3.8	Kerala 54.1 Karnataka 37.5 Tamil Nadu 7.0
Cardamom	91.5	1.9	32	133.2	Nil	Kerala 70.7 Karnataka 19.0 Tamil Nadu 10.3
Pepper	121.7	28.2	231	344.8	Nil	Kerala 96.5

Source: K.V.A. Bavappa¹.

TABLE 2: Nutrient content of the leaves of young coconut palms under NPK fertiliser experiment at CPCRI, Kasaragod

N		P ₂ O ₅		K ₂ O	
Treatment	% Content	Treatment	% Content	Treatment	% Content
500 g	1.60	500 g	0.10	750	0.55
750 g	1.70	1000 g	0.11	1500	0.80
1000 g	1.80	1500 g	0.13	2250	0.90

Source: CPCRI, Kasaragod⁴.

reddish brown. In a case of acute deficiency, the entire leaf turns orange yellow in colour and the leaflets start drying up from the tip. The frond size gets reduced.

Pillai and Davis⁵ estimated the annual removal of nutrients by middle aged bearing palms of low yield group (25 nuts/palm) at 55 kg N, 27 kg P and 84 kg K/ha. The nutrient removal by palms yielding around 60 nuts works out to 720g N, 330 g P₂O₅ and 1080 g K₂O/palm/year.

Seedlings, produced from seednuts collected from palms manured with K, displayed better vigour and growth than those obtained from unmanured palms⁶. It has been reported that in the palms well supplied with K, the first fruits were observed 5 years after planting, while it took 8 years to fruit for the palms that did not receive K⁷. In the NPK factorial experiment on the Tall variety, in sandy loam soil at CPCRI, Kasaragod, the palms to flower first were those that received 1 kg N and 1.5 kg K₂O/palm/year. Palms which suffered from K deficiency during the prebearing period remained on an average 15% less productive than those that never suffered.

In the fertiliser demonstration trials, conducted all over the west coast of India, John and Jacob⁸ observed that an application of 0.34 kg N+0.34 kg P₂O₅+0.68 kg K₂O/palm/year resulted in an increase of 35% in nut production and 44% in copra production over the cultivators' practice. Where they failed to obtain response to this application, significant yield increase was recorded when the amount of K₂O was raised to 0.90 kg/palm. Muliyar and Nelliatt⁹ observed yield response to potassium applica-

tion from the fifth year onwards in a 3⁸NPK experiment conducted at CPCRI, Kasaragod on middle aged palms in a sandy loam soil. They also found that for palms yielding less than 60 nuts annually, the optimum amount of N application ranged between 0.40 and 0.65 kg and that of potassium ranged between 0.89 and 1.21 kg/year/palm. From the results of a field experiment, with two amounts each of N, P and K, Marar and Pandalar¹⁰ concluded that the responses to N and K were equal and additive.

Muliyar and Nelli⁹ reported that nitrogen adversely affected the nut characters, viz., weight and volume of nut and copra weight per nut and that these characters were markedly improved by K nutrition. Although N application increased nut production by 16% the copra out-turn per palm increased only by 6%, because of the adverse effects. Potassium application increased nut production by 12% and copra out-turn per palm by 22%.

The amount of fertilisers recommended for the ordinary tall variety of coconut is 500 g N + 320 g P₂O₅ + 1200 g K₂O and that for high yielding genotypes 1000 g N + 500 g P₂O₅ + 1500 g K₂O/palm/year. This works out to 210 kg and 262 kg K₂O/ha/year.

ARECANUT (ARECA CATECHU)

Arecanuts grow well on laterite soils of west coast, red loams of Mettupalayam (Tamil Nadu) and alluvial soils of West Bengal and Assam. The laterite soils which have the largest area under the crop, are slightly acidic and low in general fertility¹¹. The soils are low to medium in K.

Deficiency of K causes the leaves to become rough and puckered¹². The leaf margins curl downwards. Dead tissues occurring around the margins and between leaf veins are a sure indication of K deficiency.

It is estimated that bearing arecanut palms remove 67-90 kg N, 22-34 kg P₂O₅ and 67 to 90 kg K₂O/ha/year¹³.

Lakshmanachar *et al*¹⁴ reported the results of simple manurial trials conducted in cultivators' gardens. The fertiliser amounts were 57 and 113 kg N, 45 and 91 kg P₂O₅ and 85 and 170 kg K₂O/ha/year. The treatments found economical were 170 kg K₂O/ha for the sub-montane areas and 85.0 kg K₂O/ha for the coastal regions with 57 kg N + 45 kg P₂O₅ for both regions. A 3⁴ factorial experiment to determine the fertiliser requirement of

arecanut is in progress at 5 centres under the CPCRI. The amounts of potassium being tried are 0, 140 and 280 g/palm/year. The results show that application of potash significantly reduces leaf fall and increases the weight of nuts¹⁵.

The amounts of fertilisers recommended for arecanut are 100 g N, 40 g P₂O₅ and 140 g K₂O/palm/year. This works out to 175 kg K₂O/ha.

RUBBER (HEVEA BRASILIENSIS)

Rubber thrives in well-drained fertile loamy soils. A tonne of dry latex removes 6-8.7 kg N, 1.7-2.7 kg P₂O₅ and 5.7 to 8.4 kg K₂O from the soil.

The first symptom of K deficiency in rubber plant is the mild diffused marginal paleness of the lamina. Subsequently, the marginal border becomes yellower and the mottled areas coalesce and become necrotic, at first near the tip of the leaf. In the young trees the symptoms occur in the lower canopy, while in the older trees these are observed in the upper canopy. Reduction in leaf size and bark thickness follow.

The critical level of K in the leaves ranges between 1.0 and 1.5% on drymatter basis, the leaves exposed to the sun have lower K content than that in the leaves under shade. Potassium has a marked antagonistic effect on Ca and Mg; excess K in the soil causes Mg deficiency.

Application of potassium in amounts of 50 and 100 kg/ha increases the latex yield. The increase is more marked with the larger amount and is more persistent. In a long term experiment, du Plessix *et al*¹⁶ found that trees grown without potash fertilisation took 4 and 7 years to attain critical level of leaf K when fertilised at the rate of 300 and 150 g K₂O/tree/year respectively. They also found that the K fertilised trees yielded 19 and 18% more latex than those not receiving K fertiliser.

The use of stimulants like ethrel necessitates the application of adequate quantities of potash to ensure sustained and satisfactory flow of latex. This is because of the higher nutrient content in the latex under stimulation (Table 3). The response to stimulants was absent when K fertilisation was not practised¹⁶. The latex flow increased with increasing amounts of Potash and the latex flow reduced drastically after 3 years of stopping K fertilisation.

TABLE 3: Nutrient removal by the rubber

Treatment	Yield
No stimulation	
Ethrel	

Potty *et al.*¹⁷ found that the optimum nursery was 500, 250 and 100 kg/ha maintaining maximum number of vigorous trees on the basis of various field experiments. 40 kg N, 40 kg P₂O₅ and 16 kg K₂O, 30 kg P₂O₅ and 30 kg K₂O/ha for

TEA (TEA)

Tea grows on a wide variety of soils from the Brahmaputra valley to the laterite loams of South India. A tea bush grows well on soils with pH ranging from 4.5 to 6.5.

Tea being a leaf crop, initially the tea flush contains 4-5% N and 16-20% K. In processed tea, 160-200 kg N and 60-80 kg K₂O/ha are removed. The estimated quantity of nutrients removed is 160 kg N, 9 kg P₂O₅ and 40 kg K₂O/ha.

Potassium is now considered essential to preserve the health and vigour of tea bushes and the sustained productivity of tea bushes. Deficiency causes defoliation and reduces up resistance to a number of fungi and reduces flavour and taste of tea.

In the long term fertilizer experiment, the tea yield started declining after receiving 135 kg N continuously for

under the CPCRI. The amounts of 1280 g/palm/year. The results show that it reduces leaf fall and increases the

recommended for arccanut are 100 g N, 40 g P₂O₅ and 175 kg K₂O/ha.

TEA (THEA SINENSIS)

fertile loamy soils. A tonne of dry leaves contains 1.0 to 1.5% N, 0.5 to 0.8% P₂O₅ and 5.7 to 8.4 kg K₂O from

deficiency in rubber plant is the mild diffused chlorosis. Subsequently, the marginal border becomes necrotic and coalesce and become necrotic, at first on young trees the symptoms occur in the upper part of the trees these are observed in the upper part of the bark thickness follow.

leaves ranges between 1.0 and 1.5% on leaves exposed to the sun have lower K content than those in shade. Potassium has a marked antagonistic effect on Magnesium causes Mg deficiency.

Amounts of 50 and 100 kg/ha increases yield. The trees marked with the larger amount and in the experiment, du Plessix *et al*¹⁶ found that it took 4 and 7 years to attain critical rate of 300 and 150 g K₂O/tree/year. The K fertilised trees yielded 19 and 25% more yield than the unfertilised trees using K fertiliser.

It necessitates the application of adequate amount of K to sustain a sustained and satisfactory flow of latex. The nutrient content in the latex under continuous stimulation was absent when K fertiliser was applied. Latex flow increased with increasing K fertiliser and reduced drastically after 3 years of

TABLE 3: Nutrient removal by the rubber latex with and without stimulation

Treatment	Yield	Nutrient removal		
		N	P ₂ O ₅	K ₂ O
			(kg/ha)	
No stimulation	1390	9.4	5.3	10.0
Ethrel	2570	23.9	16.5	26.8

Potty *et al.*¹⁷ found that the optimum requirement of NPK for seedling nursery was 500, 250 and 100 kg/ha N, P₂O₅ and K₂O respectively for obtaining maximum number of vigorous, healthy and buddable seedlings. On the basis of various field experiments they suggested a fertiliser schedule of 40 kg N, 40 kg P₂O₅ and 16 kg K₂O/ha for immature rubber and 30 kg N, 30 kg P₂O₅ and 30 kg K₂O/ha for mature rubber.

TEA (THEA SINENSIS)

Tea grows on a wide variety of soils, ranging from the river alluvium of the Brahmaputra valley to the Bheel soils of the Kangra Valley and the laterite loams of South India. Although tea is known as calcifuge, it grows well on soils with pH ranging from 4 to 6.

Tea being a leaf crop, initially only N was felt necessary. However, the tea flush contains 4-5% N and 1.5 to 2.0% K. For every 1000 kg of processed tea, 160-200 kg N and 60-80 kg K₂O are removed from the soil. The estimated quantity of nutrients removed by an average tea crop is 91 kg N, 9 kg P₂O₅ and 40 kg K₂O/ha/year.

Potassium is now considered necessary to obtain sustained yield and preserve the health and vigour of the bush. It is the limiting element in the sustained productivity of tea bushes, especially under Indian conditions. Deficiency causes defoliation and debilitation of bushes. It helps to build up resistance to a number of fungal diseases. Potassium also enhances the flavour and taste of tea.

In the long term fertilizer experiments at Toklai Experimental Station, the tea yield started declining after 20-25 years¹⁸. The yield from plots receiving 135 kg N continuously for 30 years was lower by 268 kg/ha than

that from plots receiving 45 kg N. In the initial stages of the experiment 135 kg N/ha gave higher yields. When K fertilisation was super-imposed, the yield went up in proportion to the K applied. An application of 67 kg K_2O /ha restored the yield level in 135 kg N/ha plots to the same level as in 45 kg N/ha continuous applications plots.

Heavy use of N, which is essential for tea cultivation, and the low pH of soil result in the depletion of K¹⁹. The effect of K nutrition becomes dominant as the age of bushes advance. Ranganathan²⁰ estimated that the average N and K requirement of tea is 170-190 kg N and 100 kg K_2O /ha. He felt that there was a definite response to increasing amounts of K application in heavy rainfall areas up to 200 kg K_2O /ha.

COFFEE (COFFEA ARABICA)

Coffee prefers deep well drained soils, slightly acidic in reaction and rich in organic matter. The striking potash deficiency symptom is the blackening of the leaf tip and leaf edges. It also causes die back and often the whole branch dries up. The berries also develop brown colour.

Coffee, being a perennial crop, continuously produces berries and fresh wood for the succeeding crop. The crop requirement of K is high during the development of coffee berries and is maximum during their ripening. The nutrient removal by the berries in the two important cultivated varieties of coffee has been estimated at 34 kg N, 5 kg P_2O_5 and 45 kg K_2O /ha in arabica and 35 kg N, 7 kg P_2O_5 and 89 kg K_2O /ha in robusta. The nutrients required for building up of the framework are twice the quantity of nutrients removed by the berries. Iyengar and Awatramani²¹ found that a mature coffee plant had 2.5% N, 0.25% P and 3% K on dry matter basis.

Mathew *et al.*²² suggested that the most efficient proportion of N : K in terms of added fertilisers is 1 : 1 excepting in cases where the amount of available potash in the soil was low. The fertiliser recommendations for coffee are 160, 120, and 160 kg NPK/ha for arabica and 120, 90 and 120 kg NPK/ha for robusta.

CASHEW (ANACARDIUM OCCIDENTALE LINN.)

Cashew can grow on sandy soils, laterites and red loams but not on heavy clay soils. Its cultivation is generally confined to inferior land unsuitable for other more remunerative crops.

A cashew tree which yields 25 kg P_2O_5 and 1.0 kg K_2O /year⁴.

Manurial experiments carried out have shown that bearing cashew trees respond significantly to balanced NPK fertilisation over cattle manure²³. NPK fertilisation of 4 kg/tree in 4 years in an established orchard of 660 g N, 266 g P_2O_5 and 580 g K_2O /tree/year²⁴ significantly reduced the prebearing period and increased the yield of 5.68 kg nuts/tree. The mean yield increased by 7 per cent by the application of N and 10 per cent by potassium application @ 580 g K_2O /tree/year by 42%.

The amount of fertilisers generally recommended is 160 kg P_2O_5 and 120 g K_2O /tree/year.

CARDAMOM (ELEPHANTHERRIS)

Cardamom, known as the "Queen of Spices", grows in the ghats at elevations of 900 m above sea level. It requires rich organic matter and available K and P. According to Aiyar²⁵, the pH of the soil should be 5.5. The available K is about 12.5 mg/kg soil. Deficiency causes poor root and shoot growth and the plant is adversely affected. Browning of leaves occurs upwards. In acute cases, the whole leaf turns brown and production of new suckers ceases, the plant dies.

The nutrient removal by cardamom is 160 kg N, 4 kg P_2O_5 and 52 kg K_2O /year. It has also been found that 1 kg cardamom requires 1 kg P_2O_5 and 200 g K_2O . The data on nutrient removal by cardamom is twice that of nitrogen.

Since cardamom produces suckers and panicles and development of panicles takes 6-8 months, there is a steady utilisation of nutrients. Much information is, however, available on cardamom in general and potash in particular. It is not available thus far from the field experiments.

the recommended dose of fertilisers is 30 g N, 10 g P_2O_5 and 60 g K_2O per clump per year. Considering the factors affecting the availability of nutrients in the soil, the Cardamom Research Station, Mudigere recommended a fertiliser dose of 75 kg N, 75 kg P_2O_5 and 150 kg K_2O /ha for a plantation yielding 100 kg dry capsules²⁷.

OIL PALM (EALIES GUINENSIS)

Oil palm is generally grown on well drained deep loamy soil, rich in humus. Potash deficiency causes development of orange spots on the pinnae of older leaves which later turn purple in colour. The pinnae become narrow with marginal and tip scorching. The dessicated and dried old leaves are retained erect on the crown.

The fertiliser recommendation is 1400 g K_2O /palm/year, the N:K ratio being 1:2. The critical concentration of K in the leaf tissue is 1.0-1.2 per cent on dry matter basis.

PEPPER (PIPER NIGRUM)

Pepper, the king of spices, thrives best in virgin humus rich soils. It is cultivated widely on red loams and sandy loams on the hill slopes of west coast. The recommended amount of fertilisers for the perennial vine is 200 g N and 60 g K_2O per standard.

EVALUATION

Although the need to apply balanced fertilisers to maintain optimum nutrient status of the soil and to obtain sustained high productivity in plantation crops is established, the acceptance by the farmers depends on the profitability of the practice. Available experimental evidences suggest N:K ratios ranging from 0.5 to 1.0. It is also known that the damages done to these perennial crops during the pre-productive stage by K deficiency may not be fully repaired by subsequent supply of adequate quantities of fertilisers²⁸. There is a long time lag between the application of inputs and the receipt of returns through increased yields. For a fair evaluation of a practice and its economics the mean response of four years or more has to be considered because of the possible variation in response caused by seasonal conditions.

The distribution and consumption of K fertilisers in the different states during the past two years indicates an awareness among the farmers

about the importance of potassium the statewise consumption ratio of 13.5 in the North and 11.0 in the largest area under plantation crop cropped area, consumed in 1% fertilisers and 6.3 per cent of the total ever, majority of the farmers still follow old practices.

FUTURE

The exact role of potassium has to be studied in depth. Since of soil types, the mobility, fixation should be examined for different situations.

ACKNOWLEDGEMENTS

Thanks are due to Dr. N.M. Crops Research Institute, Kasaragod for the preparation of the manuscript.

1. Bavappa, K.V.A. 1977. Plantation
2. Shankara Narayanan, M.P. 1974. Report of the All India Summer Institute on plantation crops : 28 & 29.
3. Pillai, N.G. 1974. Nutrition of coconut. Institute on improvement and management of coconut crops.
4. Anonymous, 1973. Annual Report of the All India Summer Institute on plantation crops.
5. Pillai, N.G. & T.A. Davis, 1963. Experimental a preliminary study. Indian Coconut Journal (supplement) : 70-80.
6. Nelliatt, E.V. 1973. NPK Nutrition of coconut (supplement) : 70-80.
7. Salgado, M.L.N. 1952. Potash and coconut. Cocon. 3(4) : 191-192.
8. John, C.M. & K. Jacob, 1959. Fertiliser application a review. First Conference of Coconut Growers, Ernakulam.
9. Muliyyar, M.K. & E.V. Nelliatt, 1971. Fertiliser application on the West Coast of India.

30 g N, 10 g P₂O₅ and 60 g K₂O per ha. Factors affecting the availability of nutrients at the Research Station, Mudigere recommended 75 kg P₂O₅ and 150 kg K₂O/ha for a coconut palm²⁷.

(NELLIAT GUINENSIS)

It is grown on well drained deep loamy soil, rich in organic matter. Development of orange spots on the leaves and they turn purple in colour. The pinnae show signs of scorching. The dessicated and dried leaves turn brown.

At 1400 g K₂O/palm/year, the N:K ratio in the leaf tissue is 1.0.

(COCOS NIGRUM)

It is best in virgin humus rich soils. It is grown on sandy loams on the hill slopes of coconut palm. Application of fertilisers for the perennial vine

PLANTATION

Application of fertilisers to maintain optimum yields and sustained high productivity in coconut palm plantation by the farmers depends on several factors. Available experimental evidences suggest that the damage to the palm is also known that the damages to the palm during the pre-productive stage by K deficiency can be corrected by subsequent supply of adequate potassium. This time lag between the application of potassium and the increase in yields through increased yields. For a fair comparison the mean response of four years of the possible variation in response

Application of K fertilisers in the different coconut palm plantation creates an awareness among the farmers

about the importance of potassium nutrition of plantation crops. Whereas the statewise consumption ratio of N to K is 5.1 in the South Zone, it is 13.5 in the North and 11.0 in the Central Zone. Kerala which has the largest area under plantation crops with only 1.8 per cent of the country's cropped area, consumed in 1976-77, 1.4 per cent of the nitrogenous fertilisers and 6.3 per cent of the potash fertilisers on all-India basis. However, majority of the farmers still do not adopt the recommended fertiliser practices.

FUTURE LINE OF WORK

The exact role of potassium in the nutrition of plantation crops with a view to determine the economic optimum N:K ratio in different soils has to be studied in depth. Since plantation crops grow on a wide range of soil types, the mobility, fixation, losses and availability of potassium should be examined for different soils and climatic situations.

ACKNOWLEDGEMENT

Thanks are due to Dr. N.M. Nayar, Director, Central Plantation Crops Research Institute, Kasaragod for help and encouragement in preparation of the manuscript.

REFERENCES

1. Bavappa, K.V.A. 1977. Plantation crops industry. *J. Plant crops* 5(1) : 1-10.
2. Shankara Narayanan, M.P. 1974. Major Coconut growing soils of India. Report of the All India Summer Institute on improvement and management of plantation crops : 28 & 29.
3. Pillai, N.G. 1974. Nutrition of coconut palm. Report of the All India Summer Institute on improvement and management of plantation crops : 34-37.
4. Anonymous, 1973. Annual Report of CPCRI, Kasaragod, 1972.
5. Pillai, N.G. & T.A. Davis, 1963. Exhaust of macronutrients by the coconut palm—a preliminary study. *Indian Cocon. J.* 16(2) : 81-87.
6. Nelliatt, E.V. 1973. NPK Nutrition of coconut palm—a review. *J. Plant. Crops.* 1 (supplement) : 70-80.
7. Salgado, M.L.N. 1952. Potash and its importance in coconut manuring. *Ceylon Cocon.* 3(4) : 191-192.
8. John, C.M. & K. Jacob, 1959. Fertiliser demonstrations on coconut in West Coast—a review. First Conference of Coconut Research Workers in India. I.C. Coc. C. Ernakulam.
9. Muliya, M.K. & E.V. Nelliatt, 1971. Response of coconut palm to N, P and K fertiliser application on the West Coast of India. *Oleagineux* 26(11) : 681-691.

10. Marar, M.M.K. & K.M. Pandalai, 1961. A comparison of chilean nitrate with ammonium sulphate as a nitrogen source for the coconut palm. *Indian Cocon. J.* 14(3) : 110-122.
11. Bhat, K.S. 1974. Establishing arecanut plantation, nutrition and fertiliser practices. Report of the All India Summer Institute on improvement and management of plantation crops : 97-99.
12. Yadava, R.B.R., K. Vellaichamy & C.K. Mathai, 1972. Role of nutrient elements and their deficiency symptoms with reference to arecanut. *Arecanut and spices Bull.* 3(4) : 4-7.
13. Anonymous, 1970. Annual Report of the Central and Regional Arecanut Research Station, Vittal for 1968. 41-45.
14. Lakshmanachar, M.S., K.G. Biddappa & T.T. Paulose, 1966. Simple manurial trials on arecanuts in ryots' gardens. *Agric. situation India*, 21 : 441-450.
15. Anonymous, 1976. Annual Report of CPCRI, Kasaragod, 1975. 87-89.
16. du Plessix, C.J., J.M. Eschbach & A. Cornier, 1973. quoted in IPI Research Topics No. 1 Fertiliser experiments, the need for long term trials.
17. Potty, S.N., M.A. Kalam, K.I. Punnose & C.M. George, 1976. Response of Hevea to fertiliser application in relation to soil fertility characteristics. *Rubb. Board Bull.* 13(3): 48-54.
18. Dey, S.K. 1971: Changes in soil fertility under continuous cropping and manuring in tea area and probable implication towards fertiliser recommendations. *Int. Symp. Soil fertility Evaluation. Vol. 1* : 843-58.
19. Kanwar, J.S. 1974. Assessment of potassium fertilisation in the tropics and subtropics of Asia—Proceedings of the 10th Congress of the IPI in Budapest : 1-23.
20. Ranganathan, V. 1973. Efficient use of fertilisers in tea under conditions of shortage for fertiliser use under limited availability. FAI group discussion on strategy (Abstract, Fertiliser news, Nov. 1973).
21. Iyengar, B.R.V. and N.A. Awathramani, 1975. Fertiliser use in Coffee. *Planters' Chronicle* 70(1) : 3-7.
22. Mathew, P.K., M.B. Rao and A. Mir. 1971, Basis of fertiliser recommendations in coffee. *Indian coffee* 35(1): 19-20.
23. Rao, V.M.M. 1974. Establishment and management of cashew plantations. Report of the All India Summer Institute on improvement and management of plantation crops. 123-27.
24. Anonymous, 1976 (b). Report of the Committee on Cashew cultivation ICAR, New Delhi : 19-27.
25. Aiyer, A.K.Y.N. 1966. Field crops of India. VIth Edn. Bangalore Print. and Pub. Co. Ltd., Bangalore.
26. Kololgi, S.D. 1974. Plantation management of cardamom. Report of the All India Summer Institute on improvement and management of plantation crops : 178-182.
27. Anonymous, 1976 (c). Cardamom in Karnataka. Published by University of Agricultural Sciences, Hebbal, Bangalore : 8-10.
28. Nelliatt, E.V. 1974. Fertiliser application to coconut. Report of the All India Summer Institute on improvement and management of plantation crops : 38-41.