

Development of pomegranate juice blended tender coconut water beverage

O. P. Chauhan*, Asha Singh, P.S. Raju and A.S. Bawa

*Defence Food Research Laboratory, Siddarthnagar, Mysore, India 570 001

Abstract

Pomegranate juice blended tender coconut water ready-to-serve beverage was developed. Response surface methodology was employed to optimize the levels of dependent variables (levels of tender coconut water, pomegranate juice, TSS and citric acid concentration). The responses studied were sensory (colour, flavour, consistency, mouthfeel and overall acceptability), pH and objective CIE colour values (L^* , a^* , b^*). A Central Composite Rotatable Design (CCRD) was used with six experiments at the central point. The data obtained were analysed employing multiple regression technique to generate suitable mathematical models. Quadratic polynomial models were found to fit well in describing the effect of variables on the responses with high correlation coefficients (R^2 87.62- 99.76), showing the adequacy of the models. An optimum level of the dependent variables (tender coconut water 6.54%, pomegranate juice 6.97%, TSS 12.8^o Brix and citric acid 0.12% was achieved through response surface models for getting optimum values of the responses. The experimental and predicted values at optimum conditions showed high correlation coefficients. The beverage prepared

at optimum conditions of the variables showed a shelf life of 6 months at 5, 25 and 35°C under packed condition.

Introduction

Tender coconut water is a very popular drink in the world market, especially for its healing qualities such as oral or intravenous rehydration. It is the natural isotonic beverage with almost the same level of electrolytes as we have in our blood (Suresh *et al.* 1968) and does not destroy red blood cells and is readily accepted by the body (Suresh and Hegde, 1971). It is very helpful in relieving cholera diseases combating intestinal worms and relieving stomach troubles (Jasper, 1979). It is the most nutritious wholesome beverage that the nature has provided. The water of fresh tender coconut has been usually drunk by people with no ill effects. Tender coconut water is not only a thirst quenching but also a mineral drink that cures most of the diseases. Tender coconut water is considered as 'unctuous, sweet, increases semen production, clears the urinary path and strengthen the muscle of cardiovascular system". Sports drinks based on tender coconut water have ample market potential (Anon, 2000). Health drinks based on coconut water are also very popular

Tender coconut water is a very popular drink in the world market, especially for its healing qualities such as oral or intravenous rehydration. It is the natural isotonic beverage with almost the same level of electrolytes as we have in our blood and does not destroy red blood cells and is readily accepted by the body.

in different parts of the world (Magda, 1992). It is good for feeding infants suffering from gastro-intestinal diseases (Nurasid *et al.* 1979, Munir and Mustadjab 1980). It also checks urinary infections, has diuretic property and effective in the treatment of kidney, urethral stones and also cures malnourishment (Macalalag and Macalalag, 1987). It is also reported to contain substances capable of inducing rapid proliferation of plant tissue and it is also used as a culture medium for microorganisms.

Coconut water is the liquid endosperm that fills the central cavity enclosed by a solid endosperm. The coconut water, technically the endosperm, is formed in small quantities in the third month of the development of the nut and reaches maximum in the 8th month and declines thereafter as the nut ripens. Its major chemical constituents are sugars and minerals while fat and nitrogenous substances form a minor fraction (Shivashankar, 1991). A drastic change in chemical composition of tender coconut water has been reported by Jayalekshmi *et al.*, (1986). As the coconut reaches towards its maturity, fat and protein content increases markedly and water, total solids, sugars, ash and mineral constituents decreases on per unit basis.

Plain tender coconut is available in the market in preserved and packaged form, but it is not very much acceptable by the consumers because of its bland taste and flavour. Therefore, an attempt was made to develop a refreshing and more acceptable ready-to-serve beverage based on tender coconut water blended with pomegranate juice as such, is an important

commercial product and is used for its colour and flavour in a wide range of juices beverages and other food products (Alper and Acar, 2004). It contains considerable amount of sugars, acids, vitamins, polyphenols and important minerals (Vardin and Fenercioglu, 2003). It is also reported to contain estrogen like activity (Maru *et al.* 2001). It is also useful in reducing oxidative stress, atherogenic modification to LDL and platelet aggregation (Aviram *et al.*, 2000).

Therefore, the present study was undertaken to develop a pomegranate juice blended tender coconut water beverage having good consumer acceptability. The levels of the tender coconut water and pomegranate juice were optimized using response surface methodology. The total soluble solids and acidity of the beverage were also optimized as they play major role in the acceptability of the RTS beverages.

Materials and methods

Fruits

Fully mature lemon and tender (green) coconuts were procured

from the local Mysore (India) market for the study. The fruits were sorted and graded and those with visible infections and mechanical injuries were rejected. The selected fruits of lemons were subjected to surface sanitation wash in chlorinated water (50 ppm). The juice was extracted manually from the lemons by cutting into halves and squeezing. Tender coconut water was taken out from the nuts by penetrating a needle in the nut and sucking the water using a vacuum machine.

Experimental design

Response surface methodology was used for designing the experiment (Khuri and Cornell 1987) to develop pomegranate juice blended tender coconut water RTS beverage. A Central Composite Rotatable Design (CCRD) was used taking 4 variables at 5 levels each with 6 points at centre. The data pertaining to the coded and uncoded parameter values of the design have been presented in Tables 1 and 2 under which 6 replications at the central point and 2 at the other points were carried out. The effect of four independent variables (pomegranate

Table 1. Process variable and their coded and uncoded levels

Factors	Process variable	-2.00	-1.00	0.00	+1.00	+2.00
A	Tender coconut water (%)	2.5	5	7.5	10	12.5
B	Pomegranate juice (%)	2.5	5	7.5	10	12.5
C	TSS (^o Brix)	5	10	15	20	25
D	Citric acid (%)	0.025	0.05	0.125	0.2	0.275

Table 2. Response surface design showing number of experiments at different levels of variables

A	B	C	D	Runs
0	0	0	0	6
±2.0	0	0	0	2
±1	±1	±1	±1	16
0	+2.0	0	0	2
0	0	+2.0	0	2
0	0	0	+2.0	2



juice, tender coconut water, TSS and citric acid concentration) were investigated on dependent responses (sensory colour, flavour, consistency, mouthfeel and overall acceptability as well as pH, and CIE L*, a* and b* values.

Optimization

The data obtained for pH, CIE L*, a*, b* values and sensory attributes (colour, flavour, consistency, mouthfeel and overall acceptability) under different experimental conditions were analyzed using multiple regression analyses. Response surface model was developed to represent the data thus collected and to determine the optimal conditions for various parameters. A quadratic polynomial regression model was assumed for predicting the response variable (Y) and the data were fitted to the following second order polynomial equation :

$$Y = \beta_0 + \sum_{i=1}^4 \beta_i X_i + \sum_{i=1}^4 \beta_{ii} X_i^2 + \sum_{i=1}^3 \sum_{j=i+1}^4 \beta_{ij} X_i X_j$$

Where β_0 is the value for the fixed response at the central point of the experiment, β_i , β_j , β_{ii} and β_{ij} are the linear, quadratic and cross product coefficients, respectively.

CIE color coordinates

The CIE colour values (L*, a* b*) were measured using D-65 illuminant and 10⁰ observer using a colour meter (MiniScan XE Plus, Model No. 45/O-S, Hunter Associates Laboratory, Inc., Reston, VA, USA). The equipment was calibrated using white and black standard ceramic tiles and the readings were recorded with inbuilt

software (Easy Match QC, Hunter Associates Laboratory, Inc., Reston, VA, USA).

Physico-chemical analysis

Ascorbic acid, titratable acidity, total and reducing sugars were analyzed as per the method described by Ranganna (1999). TSS of the beverage was measured using a hand refractometer (ERMA, Japan) whereas pH was determined using a microprocessor based pH meter (Century, Model CP 931, Bangalore, India). Total phenolics and flavonoids were estimated using the method described by Singleton *et al.*, (1999) and Zhishen *et al* (1999), respectively.

Sensory evaluation

The sensory evaluation of the beverage was carried out in terms of colour, flavour, consistency, mouthfeel and overall acceptability by a trained panel consisting of 10 members from the scientific staff of the laboratory with knowledge of consumer preferences using a nine point hedonic scale having a score of 9 for extreme liking and `1 for extreme disliking (Larmond, 1977). The samples were served to the panelists at 11 a.m. during each time after duly coding them using a sensory cubical illuminated with white light and maintained at 20°C.

Storage study

The ingredients were mixed as per the design for the development of the beverage. Sodium benzoate (150 ppm) was added and the beverage was packed (200 ml pack size) in laminated standi pouches

(12µ polyester, 9µ aluminium foil, 15µ nylon and 70µ CPP) and pasteurized in boiling water for 15 min. The samples were kept at 5, 25 and 35°C for 6 months and were periodically analyzed for physico-chemical, microbiological and sensory attributes.

Microbiological evaluation

The microbiological analysis of the beverage was carried out during storage for standard plate count, yeast and mold count and total coliform count according to APHA (1992) procedures.

Statistical analysis

Contour plotting was carried out to visualize the effect of variables on responses using response surface equation. A series of contour plots were developed to interpret the effect of variables on the responses investigated. A second order polynomial was fitted to the mean data values to obtain regression equations and statistical significance was examined using statistical software (Design Expert Software, version 7.1, Statease, USA). The experimental and computed values were analyzed for coefficient of determination (R²), standard error and scattered plot. The levels of independent variables were optimized using the same software considering the sensory responses at maximum values. Physicochemical data and sensory scores of the beverage during storage were analyzed statistically for analysis of variance (ANOVA) at P < 0.05 using Statistica 7 software (Stat Soft, Tulsa, OK, USA).



Results and discussion

Effect of dependent variables on sensory attributes

The sensory score for colour, flavour, consistency, mouthfeel and overall acceptability ranged from 5.80 to 7.58, 6.12 to 8.12, 6.03 to 7.68, 5.84 to 7.89 and 6.23 to 8.24; respectively at different experimental levels of the variables. The sensory colour was significantly ($p \leq 0.0001$) affected by all the four variables (Table 3 and 4). Tender coconut water was found to significantly ($p \leq 0.0001$) reduce the sensory colour scores whereas pomegranate juice, TSS and citric acid were found to improve the sensory scores for colour significantly ($p \leq 0.0001$). All the variables affected the sensory colour values at quadratic level with R^2 value of 99.69. Sensory scores for flavour, consistency mouthfeel and overall acceptability were found to be affected to a greater extent by the level of pomegranate juice (Fig. 1-5). The high R^2 values (87.62, 99.76, 99.76 and 95.04) showed adequacy of the polynomial model for the sensory flavour,

consistency, mouthfeel and overall acceptability, respectively. The flavour was significantly ($p \leq 0.0001$) affected more by the level of pomegranate juice followed by TSS, citric acid concentration, and the level of tender coconut water. Pomegranate juice also improved the consistency and mouthfeel of the beverage leading to higher sensory score for overall acceptability of the beverage. The effect of tender coconut water on flavour and overall acceptability of the beverage was found to be non significant. TSS also showed non significant effect on the overall acceptability of the beverage. The beverage prepared using 7.5% of pomegranate juice showed highest sensory scores for all the sensory responses and addition of tender coconut water beyond 7.5% decreased the sensory scores. The dependent variables affected the sensory response for flavour, consistency and overall acceptability significantly ($p \leq 0.05$) at linear level, whereas, the effect was not significant for colour and mouthfeel. The effect was also found to be not

significant at interactive levels of the variables for all the sensory responses.

Effect of variables on pH

The beverage samples showed a pH range of 3.05 to 3.45 at different levels of experimental variables. Quadratic model was found to fit well with R^2 value of 99.49 in describing the effect of all the variables on pH. Citric acid and concentration significantly ($p \leq 0.0001$) decreased the pH of the beverages, whereas, other variables had positive effect on pH value (Table 3 and 4). Tender coconut water had more pronounced effect on pH as compared to pineapple juice (Fig. 6). The sample prepared using 0.15% citric acid showed minimum pH value, whereas, those prepared using 0.075% citric acid showed maximum pH value. All the variables affected the pH of the beverages at linear, quadratic and interactive levels. A polynomial model was found to fit well in describing the effect of variables on pH of the beverage developed.

Table 3. Coefficient of second order polynomial regression model for pomegranate juice blended tender coconut water beverage

Coefficient	Color	Flavor	Consistency	Mouth feel	OAA	pH	L*	a*	b*
Intercept (β_0)	7.54	7.6	7.64	7.85	7.97	3.31	3.89	8.47	0.96
A-TCW (β_1)	-0.10	0.07	-0.03	0.024	0.04	0.085	0.24	-0.01	-0.007
B-Pomegranate juice (β_2)	0.05	0.36	0.34	0.28	0.4	0.03	-0.50	1.39	0.3
C-TSS (β_3)	0.078	0.18	0.05	0.11	0.08	0.02	-0.13	0.65	0.025
D-Citric acid (β_4)	0.15	0.095	0.18	0.14	0.21	-0.026	-0.09	0.64	0.005
AB (β_{12})	0.2	-0.05	-0.12	0.22	-0.03	-0.029	-0.015	-0.5	0.06
AC (β_{13})	0.27	0.043	0.22	0.30	0.23	0.009	-0.3	-0.25	2.44
AD (β_{14})	0.02	0.12	0.11	0.17	0.09	0.003	0.14	0.39	-0.16
BC (β_{23})	0.003	0.048	-0.13	0.008	-0.1	-0.008	-0.23	0.089	0.17
BD (β_{24})	0.039	-0.04	0.098	0.27	0.1	0.038	0.18	0.009	-0.18
CD (β_{34})	0.1	0.125	0.17	0.028	0.14	0.034	-0.09	0.37	0.16
A ² (β_{11})	-0.32	-0.18	-0.36	-0.30	-0.31	-0.006	0.2	-0.1	-0.2
B ² (β_{22})	-0.39	-0.12	-0.18	-0.26	-0.20	-0.01	0.59	-1.0	-0.35
C ² (β_{33})	-0.10	-0.32	-0.16	-0.35	-0.30	0.008	-0.17	-0.29	-0.08
D ² (β_{44})	-0.04	-0.30	-0.24	-0.26	-0.27	-0.003	0.29	0.014	0.29




Fig. 1. Colour as a function of dependent variables




Fig. 2. Flavour as a function of dependent variables




Fig. 3. Consistency as a function of dependent variables




Fig. 4. Mouthfeel as a function of dependent variables




Fig. 5. Overall acceptability as a function of dependent variables

Effect of variables on CIE L*, a*, b* values

Objective colour of the beverage was determined using a colour meter on CIE scale. The L* value which

the beverage increased (Fig. 7). The variables affected the L* value significantly ($p \leq 0.0001$) at quadratic level only. The a* value showing redness/greenness of the sample was

shows the lightness was found to decrease with increase in the level of pomegranate juice, TSS and citric acid. Tender coconut water showed a positive effect, indicating that with the increase in the concentration of tender coconut water L* value of

also highly dependent on the level of pomegranate juice, which increase the a* value (Fig. 8). The TSS and citric acid also showed positive effect on a* value, whereas, tender coconut water showed negative effect decreasing a* value with increase in its concentration. The variable affected a* value significantly ($P \leq 0.0001$) at linear and quadratic levels, whereas, the effect was not significant at interactive level. The yellowness/blueness is represented by b* value and it was found that with the increase in the concentration of pomegranate juice, the b* value increased significantly ($p \leq 0.0001$). Tender coconut water had negative effect on b* value, whereas, TSS and

Table 4. Analysis of variance data for response variables

Source	Sum of squares								
	Colour	Flavour	Consistency	Mouthfeel	OAA	pH	L* value	a* value	b* value
Model	9.31****	9.76***	9.99****	13.18****	12.55****	0.29****	24.53****	95.48****	10.46****
A-TCW	0.07****	0.11	0.017**	0.014*	0.039	0.18****	1.37*	0.0048	0.001
B-Pomegranate juice	0.24****	3.06****	2.77****	1.83****	3.80****	0.022****	5.87****	46.54****	2.18****
C-TSS	0.15****	0.79*	0.063****	0.30****	0.16	0.0057****	0.43	0.10	0.016
D-Citric Acid	0.55****	0.21	0.8****	0.45****	1.097***	0.017****	0.20	9.91****	0.0007
AB	0.64****	0.045	0.21****	0.80****	0.018	0.014****	0.0039	2.56****	0.059
AC	1.19****	0.03	0.78****	1.41****	0.82***	0.0014**	1.46*	1.03***	0.96*
AD	0.008	0.24	0.19****	0.48****	0.13	0.00015	0.32	2.418****	0.43*
BC	0.0002	0.037	0.27****	0.001	0.17	0.0011*	0.87	0.126	0.47*
BD	0.024**	0.03	0.15****	1.18****	0.17	0.023****	0.50	0.0012	0.53*
CD	0.16****	0.25	0.47****	0.013*	0.32*	0.019****	0.13	2.16****	0.43
A ²	2.87****	0.96*	2.58****	2.52****	2.73****	0.00097*	1.11*	0.257	1.12*
B ²	4.22****	0.42	0.95****	1.79****	1.16***	0.0033****	9.62****	29.06****	3.35****
C ²	0.29****	2.82****	0.75****	3.35****	2.39****	0.0017***	0.75	2.34****	0.16
D ²	0.064****	2.42***	1.58****	1.81****	1.93****	0.00032	2.30**	0.006	2.30**
Combined effect of variable									
Linear	1.01	4.17*	3.64*	2.59	5.09*	0.22****	7.87	56.55***	2.20
Interactive	2.02	0.63	2.08	3.88	1.62	0.059****	3.28	8.30	2.87
Quadratic	6.28****	4.96****	4.26****	6.70****	5.83****	0.0068****	13.38****	30.64****	5.39****
R ² (%)	99.69	87.62	99.76	99.76	95.04	99.49	88.73	99.09	88.15

**** Significant (P<0.0001), ***Significant (P<0.001), **Significant (P<0.0005), *Significant (P<0.05)

citric acid showed positive effect on b* value, but to a lesser extent as compared to pomegranate juice (Fig. 9). The variables affected the b* value significantly (p<0.0001) at quadratic level only. Polynomial model was found to fit well with the experimental CIE colour values with

high correlation coefficients (88.73, 99.09 and 88.15 for L*, a* and b*, respectively).

Optimization of process variables

The variables were optimized for getting optimal values of responses using Design Expert Software

(Version 7.1, Statease, USA). The compromised optimized levels of variables obtained for tender coconut water, pomegranate juice, TSS and citric acid were 6.54%, 6.97%, 12.8° Brix and 0.12%, respectively. A high correlation could be established between the




Fig. 6. pH as a function of dependent variables




Fig. 7. CIE L* value as a function of dependent variables




Fig. 8. CIE a* value as a function of dependent variables




Fig. 9. CIE b* value as a function of dependent variables

predicted and the actual response values under these optimized conditions (Table 5). Riaz and Elahi (1992) studied the effects of different levels of pomegranate juice (5, 10 and 20%) on physico-chemically and consumer acceptance of carbonated pomegranate beverage and found that beverage containing 10-20% pomegranate juice had maximum acceptability. In the present study some percentage of the pomegranate juice had been supplemented by tender coconut water for getting good consumer acceptability.

Storage study

Pomegranate juice blended tender coconut water RTS beverage was prepared at the optimum levels

of the independent variables and stored at 5, 25 and 35°C. Various physicochemically data obtained during storage study revealed that the prepared beverage was stable for a period of 6 months at all the three temperatures. TSS, pH, total phenolics, total flavonoids, total sugar and ascorbic acid decreased significantly ($p < 0.05$) whereas acidity and reducing sugars increased significantly ($P < 0.05$) during storage for 6 months (Table 6). The CIE L* and b* values decreased with increase in the storage duration. The a* value increased during storage showing increase in redness of the beverage. The sensory scores for colour, flavour, consistency, mouthfeel and overall acceptability decreased with

increase in the storage duration at all the three temperatures, but, the sensory scores remained within acceptable limit. The changes in various physicochemically parameters and sensory attributes were found to be more at higher temperatures. The samples stored at 5°C showed minimum changes in physicochemically parameters. Marti *et al.* (2002) also observed changes in ascorbic acid and anthocyanin content during storage of pomegranate juice at 5 and 25°C for 5 months and reported less change at 5°C. The beverage samples were found free of microbial counts during entire storage period showing microbial stability for a period of 6 months at all the three temperatures.

Conclusion

All the dependent variables (tender coconut water, pomegranate juice, TSS and citric acid concentration) affected the sensory, pH and objective colour responses at quadratic level. Polynomial model fits well in describing the effect of variables on all the responses studied. Pomegranate juice affected the sensory attributes and CIE colour coordinates (L*, a*, b*) to greater extent as compared to tender coconut

Table 5. Predicted and actual response values at optimized conditions

Responses	Predicted Value	Actual value
Colour	7.52	7.53
Flavour	7.49	7.48
Consistency	7.53	7.52
Mouth feel	7.72	7.70
OAA	7.84	7.82
pH	3.26	3.26
L* value	3.95	3.98
a* value	8.29	8.26
b* value	-0.018	-0.017



Table 6. Change in physico-chemical, sensory and microbiological parameters of pomegranate juice blended tender coconut water beverage during storage

Parameters	Initial	Pomegranate - TCW					
		RT		3 Months 5°C		6 Months 37°C	
TSS (°Brix)	12.8 ^a	12.8 ^a	12.8 ^a	12.7 ^b	12.7 ^b	12.8 ^a	12.6 ^c
pH	3.26 ^a	3.26 ^a	3.26 ^a	3.24 ^c	3.24 ^c	3.25 ^b	3.23 ^d
Acidity (% citric acid)	0.12 ^a	0.12 ^a	0.12 ^a	0.13 ^c	0.12 ^a	0.12 ^a	0.126 ^d
Total phenolics (mg/100 ml)	7.13 ^a	7.12 ^a	7.12 ^a	7.11 ^b	6.99 ^c	7.11 ^b	6.56 ^d
Total flavonoids (mg/100 ml)	1.20 ^a	1.19 ^a	1.19 ^a	1.16 ^b	1.13 ^d	1.15 ^c	1.12 ^e
Total Sugar (%)	12.52 ^a	12.52 ^a	12.52 ^a	12.50 ^b	12.5 ^a	12.51 ^a	12.50 ^b
Reducing Sugar (%)	2.35 ^a	2.41 ^c	2.37 ^b	2.47 ^e	2.40 ^c	2.36 ^b	2.45 ^d
Ascorbic acid (mg/100 ml)	2.02 ^a	1.94 ^c	1.98 ^b	1.86 ^e	1.91 ^d	1.95 ^c	1.82 ^f
CIE colour values							
L*	3.98 ^a	3.93 ^c	3.98 ^a	3.89 ^d	3.88 ^d	3.96 ^b	3.84 ^e
a*	8.26 ^a	8.27 ^b	8.26 ^a	8.28 ^b	8.29 ^c	8.28 ^b	8.30 ^d
b*	-0.017 ^a	-0.016 ^b	-0.017 ^a	-0.015 ^a	-0.015 ^c	-0.016 ^b	-0.014 ^d
Sensory attributes							
Colour	7.53 ^a	7.52 ^b	7.53 ^a	7.51 ^c	7.49 ^d	7.50 ^c	7.46 ^e
Flavour	7.48 ^a	7.46 ^b	7.47 ^a	7.44 ^c	7.42 ^d	7.45 ^b	7.42 ^d
Consistency	7.52 ^a	7.51 ^b	7.52 ^a	7.48 ^d	7.50 ^c	7.51 ^b	7.47 ^e
Mouth feel	7.70 ^a	7.68 ^b	7.70 ^a	7.64 ^c	7.64 ^c	7.68 ^b	7.62 ^d
OAA	7.82 ^a	7.80 ^b	7.81 ^a	7.78 ^d	7.77 ^c	7.79 ^c	7.76 ^f
Microbiological quality							
TPC (cfu/ml)	Nil	Nil	Nil	Nil	Nil	Nil	Nil
CC (cfu/ml)	Nil	Nil	Nil	Nil	Nil	Nil	Nil
Y & M (cfu/ml)	Nil	Nil	Nil	Nil	Nil	Nil	Nil

*Values with different superscript in the same row differ significantly (p<0.05)

water, TSS and citric acid concentration. The optimized beverage was found shell stable at 5, 25 and 35°C for a period of 6 months on the basis of physicochemically and microbiological attributes. The developed beverage showed good consumer acceptability even after 6 months of storage and the optimum levels of the variables can be made use of in commercial production of the pomegranate juice blended tender coconut water beverage.

References

Anon, 2000. New sports drink: Coconut water. *Fruit Processing* **10** (12) 491-492.
 Alper, N. and Acar, J. 2004. Removal of phenolic compounds in pomegranate

juices using ultrafiltration and lactase-ultrafiltration combinations. *Nahrung* **48** : 184-187.
 AOAC, 1990. *Official Methods of Analysis*. Association of Official Analytical Chemists, Inc. 15th Edn. Arlington, Virginia. USA, 1298.
 APHA, 1992. *Compendium of methods for the microbiological examination of foods*. 16th Edn. Speck, M.L. (Ed.) American Public Health Association, Washington, DC, 734.
 Aviram, M., Dornfeld, L., Rosenblat, M., Volkova, N., Kaplan, M., Coleman, R., Hayek, T., Presser, D. and Fuhrman, B. 2000. Pomegranate juice consumption reduces oxidative stress, atherogenic modifications to LDL., and platelet aggregation : studies in humans and in atherosclerotic apolipoprotein E-deficient mice. *American Journal of Clinical Nutrition* **71** (5) 1062-1076.

Jasper, C. W. 1979. *Coconuts : Production, Processing, Products*. 2nd Ed. Van nostrand reinhold, AVI, New York.
 Jayalekshmy, A., Arumughan, C., Narayanan, S. and Methew, A.G. 1986. Changes in the chemical composition. *J. of Food Sci. Tech.* **23** (4) : 203-207.
 Khuri, A.I., Cornell, J.A. 1987. *Response surface designs and analysis*. Marcel Dekker Inc New York, 127-145.
 Larmond, E. (1977). *Laboratory methods for sensory evaluation of foods*. Canada Department of Agriculture Publ, Ottawa, 1637.
 Macalag E. V. and Macalag A. L., 1987. Bukolysis: Young coconut water renoclysis for urinary stone dissolution. *Ind Surg* **72**:247.
 Magda, RR. 1992. Coco softdrink: Health beverage from coconut water. *Food Marketing & Technology* **6** (6) : 22-23.



Marti, N., Perez, Vicente, A., Garcia Viguera, C. 2002. Influence of storage temperature and ascorbic acid addition on pomegranate juice. *Journal of the Science of Food and Agriculture*, **82** (2) : 217-221.

Maru, I., Ohnishi, J., Yamaguchi, S., Oda, Y., Kakehi, K. and Ohta, Y. 2001. An estrogen-like activity in pomegranate juice. *Journal of the Japanese Society for Food Science and Technology* **48** (2) : 146-149.

Munir, M. and Mustadjab, I. 1980. Coconut water as an optional oral electrolyte solutions. *Peadiatr Indones* **20** : 38-46.

Nurasid, H., Ontoseno, T. and Purwodihroto S. 1979. The use of young coconut water in pediatric cholera. *Peadiatr Indones* **18** : 219-225.

Ranaganna, S. 1999. *Handbook of analysis*

and quality control for fruit and vegetable products. Tata McGraw-Hill Publishing Company Limited, New Delhi, India, 1-29, 163-164, 578-582.

Riaz, R.A., Elahi, M. 1992. Studies on the preparation and storage stability of carbonated pomegranate drink. *Pakistan Journal of Scientific and Industrial Research*, **35** (11) : 467-470.

Shivashankar, S. 1991. Biochemical changes during fruit maturation in coconut. *J. of plantation crops* **19**: 102-119.

Singleton, V.L., Orthofer, R., Lamuela, Ravento, R. 1999. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Cicoalteu reagent. *Methods Enzymol* **299** : 152-178.

Suresh, T. P., Raviverma, V. H., Setty, S. V. and Ranachar, T.R.S. 1968. Fluid

Therapy by tender coconut water in veterinary practice. *Ind Vet. J.* **48** : 829-837.

Suresh, T. P. and Hegde, V. R. 1971. Tender coconut water as a plasma volume extender II. Studies on physiological responses. *Mysore J. Agri Sci.* **5** : 410-422.

Vardin, H. and Fenerctoglu, H. 2003. Study on the development of pomegranate juice processing technology : Clarification of pomegranate juice. *Nahrung* **47** : 300-303.

Zhishen, J., Mengheng, T. and Jianming, W. 1999. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem* **64** : 555-559.

(Source : *Cord* 2008, 24 (2))

Coconut garden making

Spacing: Most of the existing coconut plantations are planted in the square system at a spacing of 7.5 m x 7.5 m. Adoption of hedge (rectangular) system of planting coconut with wider row spacing and rows oriented in east-west direction would increase light availability to intercrops (e.g. 9.0 m x 6.5 m = 170 palms/ha). This would facilitate growing annuals and perennials right from the time of planting coconuts. Triangular system of planting (equilateral triangle) accommodates 30 palms more than square system of planting, however it interferes with the cultural operations and growing of intercrops.

Preparation of pits: The size of the pit for planting of seedlings is 1 m x 1m x 1m in normal soils. However, in soils with rocky or hard substrate: large pits (1.2 to 1.5 m³) are preferred. After opening

the pit, it has to be filled with top soil to a height of 60cm. If FYM or any compost is available that also can be mixed with top soil and seedling can be planted inside the pit. In water logged areas mounds are made and seedlings are planted in the centre of the mound.

Time of planting: Planting is generally done during the beginning of monsoon season. This varies with states. In Tamil Nadu and Andhra Pradesh it is done in October-November for Kerala it is during June-September. In water logged areas planting is taken up towards the end of the monsoon.

Replanting or under planting: Replanting or under planting becomes necessary when the yield is very low due to old age, long-term neglect and continuous exposure to adverse conditions. Generally under planting is practiced where old palms are removed in stages over a period

of three to four years.

Peg mark the area to be under planted. Remove very poor yielder and those very close to the planting pits. Plant the seedlings in the usual way. The other trees are removed @ one-third each year during 2nd, 3rd and 4th year after planting. If the existing garden is irregularly spaced, remove old palms within 1 m radial distance in first year, 2 m distance in second year, 3 m distance in third year and the rest in fourth year. Generally flowering is delayed a little in under planted palms.

Care of young plantations: Mulching the base with coconut husks, dry leaves, etc.. Husk burial in planting pits. Irrigation during dry months @ 45 litres once in 4 days, Pitcher irrigation may be adopted in water scarcity areas. Shading during summer. Regular weed control.

Source : *CPCRI*