

## Review Article

# A REVIEW OF INDIAN WORK ON PHOSPHORUS NUTRITION OF COCONUT\*

H. HAMEED KHAN, C. C. BIDDAPPA and O. P. JOSHI\*\*  
*Central Plantation Crops Research Institute, Kasaragod 670 124, Kerala, India*

### INTRODUCTION

Fertilizer application to coconut though known to give good economic returns was not being practiced by farmers due to many reasons. From the growers point of view, economic management of an input reserve is of prime importance. Though increase in yields and growth has resulted due to general fertilizer application, phosphate requirement of coconut palm has always been a debating question when specific fertilizer recommendations were made based on soil and foliar analyses. Various workers (Eden, Gowar and Salgado, 1963; Child, 1974; Pillai and Davis, 1963) opined that P needs of the palm are less. A critical review on phosphate fertilization of coconut by Wahid, Kamala Devi and Haridasan (1977) did not project the details of experience gained in India adequately. The presentation here examines the various aspects of P nutrition of coconut in the country.

### THE PLANT

The coconut palm (*Cocos nucifera* L.), grows in the humid tropics of India

over an area of 1.1 million hectare accounting for an annual production of 5619 million nuts. It lacks a tap root and possesses a continuously growing terminal bud commonly known as cabbage. The cultivar West Coast Tall is widely grown in the coastal belt of Kerala, Karnataka, Goa, Maharashtra, Gujarat and interior of Tamil Nadu. Though other cultivars like East Coast Tall (Tamil Nadu, Andhra Pradesh and West Bengal), Tiptur Tall (Karnataka), Benaullim (Goa and Maharashtra), Gangabondam (Andhra Pradesh) are also grown in different geographical regions of the country, information on specific nutrient needs under varying agro-ecological conditions is lacking. The fertilizer dose of 500 g N, 320 g P<sub>2</sub>O<sub>5</sub> and 1200 g K<sub>2</sub>O palm<sup>-1</sup> year<sup>-1</sup> evolved for Kerala conditions is uniformly followed in the fertilizer programmes in the country.

Coconut palm is grown under highly heterogeneous conditions of soil ranging from littoral sand to clay in situations from poorly drained low lying marshes to well drained uplands

\* Contribution No. 391, Central Plantation Crops Research Institute, Kasaragod

\*\* Present address: Central Institute of Horticulture for Northern Plains, Lucknow-6, Uttar Pradesh, India

and hill tops and in acid to highly calcareous soils (Menon and Pandalai, 1960, Khan, Sankaranarayanan and Narayana, 1978).

#### PHOSPHORUS PROFILE IN THE PLANT

Concentration of a nutrient in a tissue depends on the physiological function for which it is needed at that site. Phosphorus contents of the leaves were always found to be higher in the younger leaves compared to older leaves though contents varied as a function of time and season maintaining the same gradient in the crown. The P contents were lower in the crown during summer irrespective of the fact that the palms were fertilized or not, indicating a minimal internal flux during the summer period (Table I).

Kamala Devi and Velayutham (1978) observed an increase in P content of nut

water upto sixth month and a gradual decrease thereafter in the three genotypes studied. However, the P content of kernel gradually increased with the increase in the highest values at full maturity (Table II).

#### PHOSPHORUS AND CROP RESPONSE

Phosphorus uptake is small (15 to 18 kg ha<sup>-1</sup>), nearly one tenth of the total uptake of potassium as well as chlorine. It is an important nutrient as it enters into the regeneration of adenosine triphosphate and phosphoproteins (Manciot, Ollagnier and Ochs, 1979). Phosphoric acid has been found to increase the girth at collar, number of leaves and rate of leaf production in seedlings (Mathew and Ramadasan, 1964). Deficiency of phosphate in soil retards growth, delays flowering and also ripening of nuts.

Table I. *Phosphorus distribution in the crown*

| Leaf rank | Fertilized palms |       |       | Unfertilized palms |       |       |
|-----------|------------------|-------|-------|--------------------|-------|-------|
|           | Jan.             | May   | Sept. | Jan.               | May   | Sept. |
| 1         | 0.138            | 0.122 | 0.149 | 0.149              | 0.121 | 0.149 |
| 6         | 0.136            | 0.120 | 0.129 | 0.140              | 0.118 | 0.141 |
| 11        | 0.135            | 0.116 | 0.127 | 0.140              | 0.112 | 0.136 |
| 16        | 0.127            | 0.108 | 0.125 | 0.133              | 0.106 | 0.127 |
| 21        | 0.121            | 0.095 | 0.116 | 0.118              | 0.096 | 0.118 |

Table II. *Changes in P content of nut water and kernel with maturity of nut in different genotypes*

| Month | P in nut water mg/l |     |     | Kernel P (%) |       |       |
|-------|---------------------|-----|-----|--------------|-------|-------|
|       | WCT                 | D×T | T×D | WCT          | D×T   | T×D   |
| 4     | 48                  | 40  | 44  | -            | -     | -     |
| 6     | 118                 | 120 | 131 | -            | -     | -     |
| 8     | 186                 | 194 | 192 | 0.070        | 0.075 | 0.072 |
| 10    | 140                 | 152 | 148 | 0.090        | 0.099 | 0.114 |
| 12    | 108                 | 97  | 103 | 0.207        | 0.232 | 0.214 |

Kunhi Muliya and Nelliya (1971) observed that application of P did not improve any of the nut characters, viz., weight and volume of unhusked nut and husked nut whereas it increased copra out-turn nut<sup>-1</sup> both at lower level (340 g P<sub>2</sub>O<sub>5</sub> palm<sup>-1</sup>) and at higher level (680 g P<sub>2</sub>O<sub>5</sub> palm<sup>-1</sup>) of annual application. However, the last three years of the experiment and combined 10 years' data showed significant effect with respect to yield. They opined that the significant difference in yield may probably be due to accumulation and slow release of fixed phosphorus from the soil. The response function was linear in shape and the optimum dose of P was found to be beyond the highest dose tried. Based on the response surface function, the optimum doses (palm<sup>-1</sup> year<sup>-1</sup>) of

N, P and K were found to be 0.47 kg N, 1.00 kg P<sub>2</sub>O<sub>5</sub> and 0.89 kg K<sub>2</sub>O.

Ramanandan (1964) observed that phosphoric acid application does not appear to influence yield. Increase in yield due to its application was little (Table III).

In an NPK experiment on young palms (started in 1965) on red sandy loam soil at Kasaragod, the response to applied P was obtained for two consecutive years (1978 and 1979). Higher levels of P at 1000 and 1500 g P<sub>2</sub>O<sub>5</sub> palm<sup>-1</sup> year<sup>-1</sup> gave significantly more nuts and influenced female flower production (Table IV). However, the response was not consistent and significant in succeeding years (Anonymous, 1982).

Table III. *Effect of graded doses of phosphorus application on yield of palms*

| Nitrogen                      |                                       | Potassium                     |                                       | Phosphorus                    |                                       |
|-------------------------------|---------------------------------------|-------------------------------|---------------------------------------|-------------------------------|---------------------------------------|
| Dose (kg tree <sup>-1</sup> ) | Mean yield (nuts palm <sup>-1</sup> ) | Dose (kg tree <sup>-1</sup> ) | Mean yield (nuts palm <sup>-1</sup> ) | Dose (kg tree <sup>-1</sup> ) | Mean yield (nuts palm <sup>-1</sup> ) |
| 0.00                          | 47.8                                  | 0.00                          | 50.3                                  | 0.00                          | 49.3                                  |
| 0.34                          | 55.1                                  | 0.34                          | 52.8                                  | 0.34                          | 53.0                                  |
| 0.68                          | 53.7                                  | 0.68                          | 53.5                                  | 0.68                          | 54.3                                  |
| C. D. at 5%                   | 3.34                                  |                               | N.S.                                  |                               | 3.28                                  |

(N, P and K were supplied through ammonium sulphate, superphosphate and muriate of potash respectively)

Table IV. *Nut yield and female flower production at different levels of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O in sandy loam soil (1977-'78)*

| Nutrient level (g palm <sup>-1</sup> year <sup>-1</sup> ) | Yield of nuts (nuts <sup>-1</sup> palm <sup>-1</sup> ) |    |      | Female flower production (No. palm <sup>-1</sup> ) |     |      |
|---|--|----|------|--|-----|------|
|   | N  | P  | K    | N  | P   | K    |
| 500   | 44   | 37 | 43   | 136  | 122 | 143  |
| 1000  | 51   | 52 | 50   | 164  | 168 | 158  |
| 1500  | 45   | 51 | 46   | 146  | 155 | 144  |
| S. E. (±)   | 3  | 3  | 3    | 13   | 13  | 13   |
| C. D. at 5%   | N.S.   | 11 | N.S. | N.S.   | 32  | N.S. |

The results of an NPK experiment (Balaramapuram, Kerala) initiated in 1964 revealed significantly high yield of 5386 nuts ha<sup>-1</sup> in the treatment P<sub>1</sub> (225 g palm<sup>-1</sup>) followed by P<sub>2</sub> (450 g palm<sup>-1</sup>) and P<sub>0</sub> (no P) with 4227 and 2709 nuts ha<sup>-1</sup>, respectively. As highest dose of P (450 g P<sub>2</sub>O<sub>5</sub> palm<sup>-1</sup>) continued to induce a depressive effect on yield of nuts, it was inferred that application of P<sub>2</sub>O<sub>5</sub> beyond 225 g palm<sup>-1</sup> year<sup>-1</sup> may not be advisable under typical red sandy loam soil conditions. The PK interaction was significant at 5 per cent only. Highest yield of 8491 nuts ha<sup>-1</sup> was obtained in P<sub>1</sub>K<sub>2</sub> combination followed by P<sub>1</sub>K<sub>1</sub> and P<sub>2</sub>K<sub>2</sub> with 7561 and 7377 nuts ha<sup>-1</sup> respectively and were on par with each other. Application of potassium at K<sub>1</sub> (450 g K<sub>2</sub>O palm<sup>-1</sup>) and K<sub>2</sub> (900 g K<sub>2</sub>O palm<sup>-1</sup>) showed a depression in yield when dose of P was increased to P<sub>2</sub> level. Similar was the response for PN interaction but was not significant (Anonymous, 1980).

#### NUTRIENT EXHAUST STUDIES

Pillai and Davis (1963) estimated that from a sandy soil of average fertility, 22.5 kg N, 12 kg P<sub>2</sub>O<sub>5</sub> and 34.2 kg K<sub>2</sub>O were annually removed by 70 palms growing in an acre and yielding 40 nuts palm<sup>-1</sup> year<sup>-1</sup>. It was indicated

that the maximum quantity of phosphorus is lost from the palms through the leaves and nuts (Table V).

Among the different parts of the palm analysed, kernel contains 0.56 per cent P<sub>2</sub>O<sub>5</sub> followed by petiole and leaf sheath (0.51%), spadix (0.31%), stem (0.26%), leaflet (0.24%), spathe (0.164%), husk (0.15%) and shell (0.15%). Considering all the above, they opined that the recommended dose of 340 g P<sub>2</sub>O<sub>5</sub> palm<sup>-1</sup> year<sup>-1</sup> appeared to be a luxury dose and the same can be reduced to 170 g without any adverse effect for atleast sandy soils of West Coast tract.

Various workers in their different studies projected the nutrient exhaust data by coconut palm (Table VI).

#### SYSTEM ANALYSIS AND P FERTILIZATION NEEDS

Considering the annual fluxes of nutrients, Nair (1979) visualised an annual storage of 41.7 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in pure coconut system compared to 44 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> in 30 year old coconut with 5 year old cocoa mixed crop system. Based on build up of P in root zone of coconut, activity of P solubilizing microorganisms (Nair and Rao, 1977) and sluggish response of palms to P

Table V. Percentage distribution of exhaust of nutrients by different parts

| Parts of the palm    | Percentage of total exhaust |                               |                  |      |      |
|----------------------|-----------------------------|-------------------------------|------------------|------|------|
|                      | N                           | P <sub>2</sub> O <sub>5</sub> | K <sub>2</sub> O | CaO  | MgO  |
| Nuts                 | 43.0                        | 40.0                          | 63.0             | 15.3 | 25.1 |
| Peduncles            | 4.2                         | 7.0                           | 12.1             | 3.3  | 11.4 |
| Spathes              | 3.5                         | 2.9                           | 2.7              | 4.5  | 4.9  |
| Leaves with stipules | 41.2                        | 45.1                          | 12.4             | 73.8 | 56.5 |
| Stem                 | 8.1                         | 5.0                           | 9.8              | 3.1  | 2.2  |

Table VI. *Nutrient exhaust by coconut palm*

| Reference               | Basis   | Nutrient removal (Kg) |                               |                  | Ratio |                               |                  |
|-------------------------|---|-----------------------|-------------------------------|------------------|-------|-------------------------------|------------------|
|                         |   | N                     | P <sub>2</sub> O <sub>5</sub> | K <sub>2</sub> O | N     | P <sub>2</sub> O <sub>5</sub> | K <sub>2</sub> O |
| Pillai (1919)           | 2000 nuts acre <sup>-1</sup> Yr <sup>-1</sup>                   | 8.2                   | 2.3                           | 17.2             | 1     | : 0.3                         | : 2.1            |
| Patel (1938)            | Recommendations for annual addition acre <sup>-1</sup>          | 10.9                  | 5.4                           | 27.2             | 1     | : 0.5                         | : 2.5            |
| Ramanandan (1977)       | a. 175 palms ha <sup>-1</sup> (nuts alone)                      | 22.0                  | 8.0                           | 71.0             | -     | -                             | -                |
|                         | b. - do -   | 21.0                  | 6.0                           | 55.0             | -     | -                             | -                |
|                         | c. - do -   | 17.0                  | 6.0                           | 52.0             | -     | -                             | -                |
| Pillai and Davis (1963) | 70 palms acre <sup>-1</sup> yielding 40 nuts palm <sup>-1</sup> | 22.5                  | 12.0                          | 34.2             | 1     | : 0.5                         | : 2.0            |

a - coastal sandy alluvium; b - red sandy loam; c - laterite

(Kunhi Muliya and Nelliya, 1971), a modification in the fertilizer schedule for coconut-cocoa mixed cropping system was suggested in which he recommended skipping of P application in alternate quarters to coconut thereby reducing the P input by 56 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> annum<sup>-1</sup> (Table VII).

#### INTERACTIONS WITH OTHER NUTRIENTS

Phosphorus and K contents of the leaf were found to be highly correlated ( $r = +0.53^{**}$  to  $0.74^{**}$ ) according to Wahid, Kamala Devi and Haridasan (1977). Our studies indicated that constant levels of N and K and varying P levels could positively influence the absorption of N ( $r = +0.47^{**}$ ) as judged by foliar level. The trend was not always significant. The increasing soil available P levels were also not found to influence either way the contents of Fe, Mn, Cu and Zn in the palm. Though a depression in the absorption of Fe, Mn and Zn was seen at higher P levels, the differences shown by the palms are not statistically significant.

#### NUTRIENT TRANSFORMATION FOLLOWING FERTILIZER APPLICATION

Kamala Devi and Velayutham (1977) found that maximum P concentration in the 14th leaf (0.170%), unopened leaf (0.146%) and outer leaf (0.139%) were obtained on the fifth day after fertilizer application. The increase in P concentration was 14-25 per cent in unopened leaf, 29-47 per cent in the 14th and outer leaves. They could not visualise appreciable differences among the genotypes (WCT, D×T and T×D) in the concentration of nutrients. For N, P and K, the best time of sampling the reference leaf to ascertain the magnitude of fertilizer effect on the nutrient content was indicated as five days and two days respectively after fertilizer application. Eventhough, younger leaves contain more phosphorus, the study indicated that the initial absorption appears to be controlled by physiological maturity of the leaves.

#### FERTILIZER PLACEMENT STUDIES

Fertilizer utilization efficiency of a crop is governed by the extent of root

Table VII. *Modifications in the fertilizer schedule for coconut + cocoa mixed cropping (kg ha<sup>-1</sup>)*

| Period     | N       |           | P <sub>2</sub> O <sub>5</sub> |           | K <sub>2</sub> O |           |
|------------|---------|-----------|-------------------------------|-----------|------------------|-----------|
|            | Present | Suggested | Present                       | Suggested | Present          | Suggested |
| February   | 43.8    | 43.8      | 28.0                          | 28.0      | 105.0            | 105.0     |
| May        | 43.8*   | 21.0      | 28.0                          | Skip.     | 105.0            | 52.5      |
|            | 17.5**  | 17.5      | 7.0                           | 7.0       | 24.5             | 24.5      |
| August     | 43.8    | 21.0      | 28.0                          | Skip.     | 105.0            | 52.5      |
| October    | 43.8*   | 43.8      | 28.0                          | 28.0      | 105.0            | 105.0     |
|            | 17.5**  | 17.5      | 7.0                           | 7.0       | 24.5             | 24.5      |
| Total      | 210.2   | 164.6     | 126.0                         | 70.0      | 469.0            | 364.0     |
| Difference |         | (-45.6)   |                               | (-56.0)   |                  | (-105.0)  |

\* Coconut; \*\* Cocoa

surface available for absorbing nutrients and the concentration of nutrients in the soil solution. Kushwah et al. (1973) exploring the rooting pattern of coconut reported the presence of only few roots in the 30 cm layer of the soil and a majority of the roots (82%) in 30-120 cm depth in regularly manured and cultivated plots. As P mobility is restricted (Barber, 1962), its placement in the most absorbing zone will be ideal for efficient utilization of nutrient. Studies employing <sup>32</sup>P on the uptake pattern of coconut (Dwivedi, Ray and Ninan, 1981) revealed that trench method of application of P is better than basin method, strip method and hole method. Eventhough <sup>32</sup>P was detected in the foliage (first fully opened leaf) after 7, 8 and 11 days in hole, trench and strip and basin method respectively, average highest accumulation of activity (cpm g<sup>-1</sup> dry matter) resulted due to trench method of application. The studies also revealed the superiority of plant injection techniques, particularly through the cut end of roots over soil placement methods; the rate of absorption of activity found by former

was 10 to 60 times faster over latter. Plant injection techniques also resulted in the detection of activity after 4, 8, 12 and 18 hours after application through cut end of roots, stem, leaf axils and growing root tips, respectively. The results of the investigation advocated that the fertilizer application through trenches around the bole of a palm for normal areas and feeding of nutrients through plant injection methods for problematic areas (water saturation and diurnal soil water fluctuations etc.) may prove ideal.

#### P CARRIERS FOR COCONUT

Khan et al. (1985) while evaluating the comparative efficiency of four phosphate carriers, viz., superphosphate, ammonium phosphate, nitro phosphate and rock phosphate, recommended rock phosphate as the ideal carrier of P for coconut as the soil contents receiving rock phosphate gave a best reflect on plant P and enriched all P fractions in the soil and influenced overall yield. Rock phosphate is also cheaper and more suitable for acid soils.

Ramanandan (1977) compared the usefulness of tri-sodium phosphate (an alkaline carrier with 17.4%  $P_2O_5$  and a byproduct obtained from extraction of rare earths from monozite sand) with single superphosphate on coconut. During 4th, 5th and 7th year, single superphosphate was found to be significantly superior to former and both were superior to control. Not much could be discerned on pH of soil due to the use of TSP. However, Khan et al. (1985) recorded significantly lower pH by the use of ammonium phosphate as P carrier.

#### FERTILIZER MANAGEMENT

Utilisation of build up reserves in soil is the most ideal and economic way of management of coconut groves. Khan et al. (1983) indicated that phosphate fertilizer application can profitably be skipped (withheld) for at least six years in situations where soil available P is around 20 to 25 ppm in 30 to 60 cm depth in coconut basin. They arrived at this conclusion after monitoring soil levels, plant levels and yield of palms. Interestingly they observed that plant P levels were not influenced even when soil available P levels drained to 59 ppm at 0 to 30 cm depth and 10 ppm at 30 to 60 cm depth (Fig. 1) and also when the soil available P values increased considerably. The yield variations were never statistically significant. The usefulness of foliar P levels as a guide to fertilizer application and yield judgement were questioned. Wahid et al. (1975) observed that when the fertilizer application was discontinued for one year,

the foliar levels of N and K were lowered and not that of P. Kamala Devi, Velayutham and Haridasan (1976) did not observe any increase in leaf P content due to graded doses of fertilizer application.


#### MYCORRHIZAL ASSOCIATION IN COCONUT NUTRITION

Most plant species growing under natural environment harbour mycorrhizae for mutual benefit. Lily (1975) first reported mycorrhizal infection in coconut (*Endogone fasciculata*) but did not relate it to phosphate nutrition. Ramesh, Joshi and Khan (1984) identified four species of mycorrhizae, namely, *Glomus mosseae* (GM), *Gigaspora gigantea* (GG), *Glomus macrocarpus* (GMA) and *Sclerocystis rubiformis* (SR) associated with coconut roots. They observed reduction in mycorrhizal infection with the increase of added as well as available P (Table VIII). Spore density of *Glomus mosseae* decreased with the addition of highest level of P. Aluminium bound P expressed as per cent of total fractions was found to be negatively related to mycorrhizal infection percentage. In view of the positive relationships between available P and spore density of GG, GMA and SR, the possibility of inoculating the coconut roots with suitable fungus to effectively recycle the accumulated P levels of soils was visualised.

#### PHOSPHORUS APPLICATION AND COCONUT DISEASES

Potty and Radhakrishnan (1978) inferred from a nutritional experiment a distinct and pronounced influence of P fertilization on stem bleeding incidence,

Fig. 1. Influence of phosphorus skipping ( $P_0$ ) and fertilization ( $P_1$ ,  $P_2$ ) on available P-PPM



$P_1=160$  g  $P_2O_5$ /palm/year;  $P_2=320$  g  $P_2O_5$ /palm/year

Phosphorus nutrition of coconut




Table VIII. Relation between different forms of P and VA mycorrhizae in coconut rhizosphere

| Treatment      | VA mycorrhizal infection percentage | population density (no. g <sup>-1</sup> ) |       |      |      | Leaf P (%) | Available P (ppm) |       | Al-P as % of total |
|----------------|-------------------------------------|---|-------|------|------|------------|-------------------|-------|--------------------|
|                |                                     | GM  | GG    | GMA  | SR   |            | Bray I            | Olsen |                    |
| M <sub>0</sub> | 82.3                                | 47.5                                      | 16.6  | 1.3  | 3.6  | 0.106      | 7.6               | 6.2   | 1.13               |
| M <sub>1</sub> | 57.3                                | 68.7                                      | 218.4 | 41.2 | 47.2 | 0.110      | 257.9             | 83.7  | 67.12              |
| M <sub>2</sub> | 42.5                                | 11.1                                      | 97.5  | 31.6 | 33.1 | 0.122      | 384.5             | 120.0 | 71.5               |

M<sub>0</sub> - No fertilizers; M<sub>1</sub> - 500 g N, 500 g P<sub>2</sub>O<sub>5</sub>, 1000 g K<sub>2</sub>O palm<sup>-1</sup> year<sup>-1</sup>; M<sub>2</sub> - 1000 g N, 1000 g P<sub>2</sub>O<sub>5</sub>, 2000 g K<sub>2</sub>O palm<sup>-1</sup> year<sup>-1</sup>

which increased with increase in the levels of P. They observed that stem bleeding was more when nitrogen supply was deficient and soil P availability was in excess. Bhaskaran, Jaganathan and Chandrasekar (1978), however, observed that higher doses of nitrogen and potash favoured faster rate of increase in disease in Thanjavur wilt of coconut, whereas P even at higher levels lowers the rate. On the contrary, Cecil et al (1978) indicated that in root (wilt) diseased coconut palms the main effect of P was not found favourable on any of the characters studied, viz., girth at collar, production of leaves, functioning leaves, flowering and yield of nuts.

CONCLUSION

Though phosphorus is a very important nutrient for coconut, the information presented suggests that phosphorus is normally not a limiting nutrient for coconut production. More so, adult palms have not been found to be much benefitted by annual P applications. Fertilizer experiments have shown that the P needs are low, responses slow and inconsistent.

Fertilizing young coconut palms upto bearing stage and experiments to

study the effect of discontinuing P application periodically thereafter will be worth investigating.

Soil P levels are to be relied more than foliar levels in judging the P needs of coconut groves. Evaluation of critical soil P levels under constant N and K fertilization over years will be rewarding in view of the lack of relationship of plant P levels on yield.

Experiments directed towards isolation and incorporation of specific species of mycorrhizae at graded levels of P application will lead to draw valuable information on the effective utilization of accumulated P reserves of the soil consequent to fertilizer application.

A survey of laterite, red sandy loam and sandy soils to study the soil and plant P levels with contemplated studies on obtaining information on requirement of P fertilization, soil critical levels and soil test methods will be other areas worth investigating.

Evaluation of coconut germplasm material to identify cultivars which show more nutrient absorption and

utilization, particularly of phosphorus under a given situation, to use them in evolving ideal genotypes for better utilization of P should also be the concern of breeders as such habits are linked with higher productivity (Beaufils, 1956).

#### ACKNOWLEDGEMENTS

The authors are thankful to Dr. K. V. Ahamed Bavappa, Director, Central Plantation Crops Research Institute, Kasaragod for the encouragement and to Dr. P. Rethinam, Project Coordinator, AICCAIP, Kasaragod for reading through the manuscript.

#### REFERENCES

- ANONYMOUS. 1982. Annual Report 1979, Central Plantation Crops Research Institute, Kasaragod, India. pp. 277.
- ANONYMOUS. 1980. Research Report 1979-'80. Directorate of Research, Kerala Agricultural University, Vellanikkara, Trichur. pp. 425.
- BARBER, S. A. 1962. A diffusion and mass flow concept of soil nutrient availability. *Soil Sci.* 93: 39-49.
- BEAUFILS, E. R. 1956. Recherche d'une exploitation rationnelle de l'Hevea basee sur l'analyse minerale de diverses parties de la plante. *Sixth Int. Congress Soil Sci.*, Paris (Colloque sur l'analyse des plantes).
- BHASKARAN, R., JAGANATHAN, T. and CHANDRASEKAR, G. 1978. Effect of fertilizers on Thanjavur wilt of coconut. In *Proc. PLACROSYM I*. 1978. pp. 351-353.
- CHILD, R. 1974. *Coconuts* 2nd ed., Longman, London, pp. 335.
- DWIVEDI, S., RAY, P. K. and SUNNY NINAN. 1981. Studies on methods of inorganic nutrient application in coconut. *Plant and Soil*. 63: 449-456.
- EDEN, T., GOWAR, T. C. and SALGADO, M. L. M. 1963. A factorial fertilizer experiment on coconuts. *Emp. J. Exp. Agric.* 31: 283-295.
- KAMALA DEVI, C. B. and VELAYUTHAM, M. 1977. Leaf nutrient concentration in coconut following fertilizer application. *The Philippine J. Coconut Studies*. 11(4): 26-33.
- KAMALA DEVI, C. B. and VELAYUTHAM, M. 1978. Changes in the chemical composition of nut water and kernel during development of coconut. In *Proc. PLACROSYM I* pp. 340-346.
- KAMALA DEVI, C. B., VELAYUTHAM, M. and HARIDASAN, M. 1976. Soil and leaf analysis in relation to nutrition of high yielding coconut genotypes. In *Coconut Research and Development*. Ed. Nayar, N. M., Wiley Eastern Ltd, pp. 518.
- KHAN HAMEED, H., SANKARANARAYANAN, M. P. and NARAYANA, K. B. 1978. Characteristics of coconut soils of India. I. Morphology, some physico-chemical characters and taxonomy. In *Proc. PLACROSYM I*: pp. 54-79.
- KHAN HAMEED, H., SANKARANARAYANAN, M. P., GEORGE, M. V. and NARAYANA, K. B. 1983. Effect of phosphorus skipping on the yield and nutrition of coconut palm (*Cocos nucifera*, L.). *J. Plant. Crops*. 11: 129-134.
- KHAN HAMEED, H., SANKARANARAYANAN, M. P., JOSHI, O. P., GEORGE, M. V. and NARAYANA, K. B. 1985. Comparative efficiency of selected phosphates as P-carriers for coconut (*Cocos nucifera* L.). *Tropical Agriculture (Trinidad)*. 62: 57-61.
- M. KUNHI MULIYAR and NELLIAT, E. V. 1971. Response of coconut palms (*Cocos nucifera* L.) to N, P and K fertilizer application on West Coast of India. *Oleagineux, Fr.* 26: 687-691.

- KUSHWAH, B. L., NELLIAT, E. V., MARCOSE, V. T. and SUNNY, A. F. 1973. Rooting pattern of coconut. *Indian J. Agron.* 18: 71-74.
- LILY, V. G. 1975. A note on development of VA mycorrhizae, *Endogone fasciculata* in coconut root. *Curr. Sci.* 44: 201.
- MANCIOT, R., OLLAGNIER, M. and OCHS, R. 1979. Nutrition mineral et fertilisation due cocotier dans le Monde. Communication presentee a la 5e session du Grupe de travail F.A.O. sur la production la protection et la triatment da la noix de coco, 3-8 December, 1979. Manila (Philippines).
- MATHEW, C. and RAMADASAN, A. 1964. Effect of NPK nutrients on the growth of coconut seedlings. *Indian Cocon. J.* 17: 114-119.
- MENON, K. P. V. and PANDALAI, K. M. 1960. *Coconut Palm*. A monograph, Indian Central Coconut Committee, Ernakulam. pp. 384.
- NAIR, P. K. R. 1979. *Intensive Multiple Cropping with Coconuts in India. Principles, Programmes and Prospects*. Verlag Paul Parey, Berlin and Hamburg. pp. 147.
- NAIR, S. K. and RAO, S. 1977. Distribution and activity of phosphate solubilizing microorganisms in the rhizosphere of coconut and coconut and cocoa under mixed cropping. *J. Plant. Crops.* 5: 67-70.
- PATEL, J. S. 1938. *The coconut - A monograph*, Govt. Press, Madras. pp. 313.
- PILLAI, N. K. 1919. *Coconut - The Wealth of Travancore*, Agri. J. India. 14: 608-625.
- PILLAI, N. G. and DAVIS, T. A. 1963. Exhaust of nutrients by the coconut palm - a preliminary study. *Indian Cocon. J.* 16: 81-87.
- POTTY, N. N. and RADHAKRISHNAN, T. C. 1978. Stem bleeding disease of coconut - nutritional relations. In *Proc. PLACROSYM I.* pp. 347-350.
- RAMANANDAN, P. L. 1964. Investigation on the nutrient inter - relationship in soils and coconut palm tissue. M. Sc. Thesis, University of Kerala, Trivandrum, Kerala.
- RAMANANDAN, P. L. 1977. Nutritional studies in coconut soils and coconut palm. Ph.D. Thesis, University of Kerala, Trivandrum, Kerala.
- RAMESH, C. R., JOSHI, O. P. and KHAN, HAMEED H. 1984. Influence of soluble phosphate on VA mycorrhiza in coconut (*Cocos nucifera* L.) - Communicated to *New Phytologist*.
- WAHID, P. A., KAMALA DEVI, C. B. and HARIDASAN, M. 1977. A critical review of phosphate fertilization of coconut. *The Philippine J. Coconut Studies.* 11(4): 1-8.
- WAHID, P. A., KAMALA DEVI, C. B., PHILIP, G. and PILLAI, N. G. 1975. Effect of discontinuation of fertilizer application on the NPK nutrition of coconut palm. *J. Plant. Crops.* 3: 58-60.